

APPLICATION FOR UNITED STATES LETTERS PATENT

5

for

COMPOSITIONS AND METHODS OF NEMATODE CONTROL

(Filed under 35 U.S.C. § 119 with Right to Priority)

sub A.7

10 This application claims benefit of U.S. Patent Application Serial No. 60/205,829 filed on May 19, 2000, entitled "Control of Nematodes, Stimulation of Nematode Resistance, and Screening Methods for Identifying Anti-Nematode Factors", incorporated herein by reference in its entirety and U.S. Patent Application Serial No. 60/274,358 filed on March 08, 2001, entitled "Control of Nematodes", incorporated herein by reference in its entirety.

15 Be it known that we, David Greenstein, a citizen of The United States of America, residing at 104 Groome Drive, Nashville, TN 37205 and Michael A. Miller, a citizen of The United States of America, residing at 2511 Barton Ave., Nashville, TN 37212; have invented new and useful "Compositions and Methods of Nematode Control".

FIELD OF THE INVENTION

20 The present invention is related to compositions and methods of nematode control and, in particular, to compositions and methods for control of nematode fertility, for identifying anti-nematode agents, and potentiating host resistance.

BACKGROUND OF THE INVENTION

Parasitic nematodes infection of plants and animals are widespread with approximately 3 billion people being infected worldwide, 100 million lives lost, and an estimated \$80 billion worth of crops lost annually to these organisms. Human conditions include river fever and elephantiasis each of which cause terrible human suffering. Parasitic nematodes are also a major problem in livestock, horses, and pets. Free living nematodes also damage plants during feeding, compete for oxygen, and transmit disease. Certain anti-nematode agents are commercially available. Disadvantages to these agents, such as the carbamates, include their extreme toxicity to most animals and all humankind. Other agents, such as ivermectin and its derivatives have undesirable toxic effects and potentially severe side effects especially those related to behavior and mental health. In addition, ivermectin resistant strains of nematodes develop relatively quickly and are reported in crops, livestock, and humans.

It is clear from the widespread and sever nature of the nematode problem and the adverse or toxic nature of many nematode treating agents, that more effective compounds and methods for controlling nematodes and identifying anti-nematode agents are needed.

STATUS OF THE PRIOR ART

McCarter et al, (1999) discloses that in the absence of sperm, the production of oocytes remains arrested in nematodes.

Klass, M. R., *et al.* (1981) discloses that major sperm protein is a structural protein in sperm cells of nematodes.

It is disclosed that all cells of *Caenorhabditis elegans* are directly observable in the intact *Caenorhabditis elegans* animal as reviewed in Hubbard and Greenstein (2000).

Video microscopy is disclosed to be used for observing the late stages of oocyte
5 development (Ward and Carrel, 1979; Albertson, 1984; Albertson and Thomson, 1993; McCarter *et al.*, 1997; Rose *et al.*, 1997; Hall *et al.*, 1999).

SUMMARY OF THE INVENTION

The present invention provides, in part, compositions and methods for controlling nematode populations, identifying anti-nematode agents, enhancing host
10 resistance to nematode infection, and treating plants and animals for nematodes.

Although the present invention is not bound by mechanism or theory, it is related, in part, to the surprising discovery made by the inventors that the major sperm protein (MSP) of nematodes is a molecular signaling factor which stimulates maturation of the female reproductive system in nematodes. Biological activities of
15 MSP in this regard include, but are not limited to: oocyte maturation, gonadal sheath cell contraction, and ovulation.

One advantage of the present invention is that inhibitors of the MSP signaling mechanism described herein are contemplated to be highly specific to inhibiting nematode proliferation and spread while being non-toxic to vertebrates
20 because the MSP gene and polypeptide are highly conserved between groups and divisions within the Phylum Nematoda, including the genera and species. Vertebrates and other non-nematode organisms, on the other hand, are not known to have an MSP gene, or protein. Thus, inhibitors of MSP stimulated female sexual maturation (FSM) are expected to be specific to organisms of the Phylum
25 Nematoda.

Another advantage of the present invention is that the high sequence conservation observed among MSP from various nematodes suggests that resistance to anti-nematode agents that target MSP is likely to be minimal as mutational changes in MSP in are likely to result in a reduction of reproductive capacity.

5 Still another advantage of the present invention is that, in general, MSP and FSM effective domains thereof, can be prepared in soluble form. Thus, MSP provides an easily handled target for methods of the present invention including in high-throughput assays for identification of MSP binding and FSM blocking agents.

10 One aspect of the present invention includes a method for identifying an anti-nematode agent by contacting a test compound to a nematode and monitoring the FSM response, wherein inhibiting test compounds are selected as anti-nematode agents.

15 Another aspect of the present invention includes identifying anti-nematode agents by selection of major sperm protein binding agents as many of these agents will also inhibit FSM. Screening methods are described to determine which MSP binding agents also inhibit FSM.

In a further aspect of the present invention, a reduction or an absence of MSP signal transduction impairs or virtually eliminates nematode fertility.

20 Still further aspects of the present invention provide methods of controlling nematode populations including, but not limited to: free living nematode populations and parasitic nematode populations which, in turn include animal and plant parasitic nematodes.

25 Additional aspects, embodiments, and elements of the present invention are described below, including in the detailed description of the invention, the examples, and the claims. Aspects, embodiments, and elements described herein are not meant to limit the present invention in any way including to any particular set

thereof. Further aspects, embodiments, elements and equivalents thereof, will be readily apparent based upon the present disclosure and are considered to be within the spirit and scope of the present invention.

BRIEF DESCRIPTION OF THE DRAWINGS

5 FIG. 1 shows a cross-section of a nematode and includes a depiction of the reproductive anatomy. Microinjections as described herein are preferred to be carried out at the point indicated.

10 FIG. 2 shows a polyacrylamide gel electrophoresis of crude sperm-conditioned medium (SCM) (left lane) and isolated 14 kDa female sexual maturation (FSM) stimulating factor (single band in right lane) identified as major sperm protein (MSP) of *C. elegans*.

FIG. 3 shows HPLC traces of fractionation of SCM on C-4 and C-18 columns. The "+" sign indicates the respective fractions with FSM positive biological activity.

15 FIG. 4 shows mass spectra of the FSM positive fractions from the HPLC purification. The mass spectra confirm MSP-3 and MSP-142 of *C. elegans* are present in the biologically active SCM. No other factors were observed or identified in these mass spectra.

20 FIG. 5 is a diagram of MSP mediated cellular communication between female reproductive cell (for example, the oocytes) and sperm.

Sub A2 21 FIG. 6 displays an alignment of twenty-seven MSP polypeptides from *C. elegans*. The SEQ ID Numbers are not meant to include the N-terminal most methionine which is believed to be cleaved during processing.

25 FIG. 7 displays and alignment of *Ascaris suum* alpha and beta MSP isoforms and MSP-142 of *C. elegans*. Again, the SEQ ID Numbers are not meant

to include to N-terminal most methionine which is believed to be cleaved during processing.

FIG. 8 is a bar graph demonstrating bioactive properties of MSP-77 and MSP-38, and sperm protein (versus and buffer control) in stimulating FSM in nematodes. The top panel displays maturations per hour which is a biological measure of oocyte maturation. The center panel displays contractions per minute which is a biological measure of sheath cell contraction. The bottom panel displays average displacement (in microns) which is another measure of sheath cell contraction. Measurements are made for buffer and the shown concentrations of MSP-77, MSP-38, and sperm MSP. The 6His marking denotes that the MSP includes a histidine tag (Qiagen) and was purified using this system.

FIG. 9 shows an alignment (performed by visual inspection) of the C-termini of MSPs from wide ranging nematode species and demonstrates that there is a remarkable sequence conservation. The residues numbers relative to the full length (minus the methionine) polypeptide are 106-126 for each of the following. *Ascaris suum* (As) MSP isoforms alpha and beta (GenBank accession numbers P27439 and P27440) fragment from 106-126 are both represented by SEQ ID NO:13 (identical polypeptides). *Onchocerca volvulus* (Ov) MSP isoforms 1 and 2 (P13262 and P13263) are also represented by SEQ ID NO:13. *Globodera rostochiensis* (Gr, the potato cyst nematode) MSP isoforms 1, 2, and 3 are each represented by SEQ ID NO:14. *C. elegans* (Ce) MSP isoforms 142 and 33 (P53017 and P53019) are each represented by SEQ ID NO:15.

FIG. 10 is a bar graph showing that the N-terminal region of MSP is necessary and sufficient for stimulation of oocyte maturation and the C-terminal region is necessary and sufficient for stimulation of sheath cell contraction. The N-terminal residues 1-106 (SEQ ID NO: 16) of MSP-77 (SEQ ID NO:9) are necessary and sufficient for stimulation of oocyte maturation (top panel). The C-terminal residues 106-126 (SEQ ID NO:17) of MSP-77 (SEQ ID NO:9) are necessary and sufficient for stimulation of sheath cell contraction (middle panel) and displacement (bottom panel).

DETAILED DESCRIPTION OF THE INVENTION

Given the human suffering and economic loss due to nematodes, it is critical that effective and safe anti-nematode compounds and methods for controlling nematodes are identified. Although not bound by mechanism or theory, the present invention takes advantage of the discovery by the inventors that nematode major sperm protein regulates the fertility of nematodes. Provided herein are compositions and methods for inhibiting or blocking major sperm protein action in fertility.

Certain utilities of the compositions and methods of the present invention include, but are not limited to: identifying anti-nematode agents, manufacturing anti-nematode agents, providing reagents for screening test compounds for anti-nematode activity, controlling nematode populations, treating animals and plants for nematode infection or infestation, treating animals and plants for certain negative effects of nematodes (including environmental or other effects of free living nematodes), raising host resistance to nematode infection, and prophylactic treatments to retard nematode infection or the spread of nematodes.

1.0 Definitions

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. In case of conflict, the present document, including
5 definitions, will control.

Descriptions of preferred methods and compositions are provided herein, but should not be construed to be limiting.

As used herein "prophylaxis" or "prophylactic treatment" refers to measures designed to preserve health or retard the spread of disease. "Prophylaxis" or
10 "prophylactic treatment" do not mean herein a certainty of the preservation of health or a certainty of a halt to a spread of a disease.

MSP is an abbreviation for major sperm protein.

FSM is an abbreviation for female sexual maturation.

FSM is meant to include, but is not necessarily limited to oocyte maturation,
15 sheath cell contraction, and ovulation.

SCM is an abbreviation for sperm-conditioned media.

C. elegans is an abbreviation for the nematode genus and species *Caenorhabditis elegans*.

A. suum is an abbreviation for the nematode genus and species *Ascaris suum*.

20 The term "biological activity" is meant to include, but is not limited to: FSM, oocyte maturation, sheath cell contraction, and ovulation.

The term "polypeptide" is known in the art, meanings of which are included herein. However, in the event of conflict, the term "polypeptide" means herein, an amino acid polymer of two units or greater (*e.g.*, a dipeptide or greater).

The term "polynucleotide" is known in the art, meanings of which are included herein. However, in the event of conflict, the term "polynucleotide" means a nucleic acid polymer of two units or greater (e.g., a dinucleotide or greater).

5 The term "isolated polypeptide" refers to a polypeptide that is at least partially removed from the milieu of molecules in which it occurs in nature.

10 The term "isolated polynucleotide" refers to a polynucleotide that is at least partially removed from the milieu of molecules in which it occurs in nature. As used herein, "isolated polynucleotide" also means that the polynucleotide is not identical in structure to a naturally occurring genome or fragment of a genome that spans more than three distinct, non-overlapping, genomically consecutive genes in length.

Additional definitions of specific terms and phrases are provided herein as needed.

2.0 *Caenorhabditis Elegans* as a Model System

15 *Caenorhabditis elegans*, or *C. elegans*, is a widely accepted genetic model system for studying the genes and gene functions of higher organisms. *C. elegans* is also a widely accepted model system for studying all features of other members of the phylum Nematoda. These features include germline development, female sexual maturation, oocyte maturation, sheath cell contraction, and ovulation (reviewed by Hubbard and Greenstein, 2000).

20 *C. elegans* is a primitive organism which nonetheless shares many of the essential biological characteristics that are central problems of, for example, human biology. The worm is conceived as a single cell which undergoes a complex process of development, starting with embryonic cleavage, proceeding through morphogenesis and growth to the adult. It has a nervous system including a rudimentary brain,
25 exhibits behaviors, and can "learn". It produces sperm and eggs, mates and

reproduces. After reproduction it gradually ages, loses vigor and finally dies. Certain genetic features of *C. elegans* have been extensively characterized and the genome has been sequenced. All 959 somatic cells of its transparent body are visible with a microscope, and its average life span (in the normal state) is a mere 2-3
5 weeks. Thus, *C. elegans* provides an ideal compromise between complexity and tractability.

3.0 Germline Development in Nematoda

In general, sexual reproduction of nematodes depends on coordination between meiotic cell cycle progression, gametogenesis, and fertilization. For
10 example, gamete differentiation is coordinated with meiotic cell cycle transitions so that fertilization produces a diploid zygote capable of completing embryogenesis and growing into a fertile adult. Normally, nematode oocytes arrest during diplotene/diakinesis of meiotic prophase (after the completion of meiosis I and meiosis II) while they grow in size. The release of oocytes from this arrest occurs
15 during meiotic maturation during which the nuclear envelope (*i.e.*, germinal vesicle) breaks down, the cytoskeleton rearranges, and the oocyte prepares for fertilization.

Typically, the progression of germline development in *C. elegans*, and members of Nematoda in general, is as follows. During embryogenesis a reproducible and largely invariant cell lineage generates two germline precursor
20 cells, Z2/Z3, and two somatic gonadal precursor cells, Z1/Z4, that together comprise the gonadal primordium at hatching (Kimble and Hirsh, 1979; Sulston *et al.*, 1983). During post-embryonic development, Z1 and Z4 give rise to the entire somatic gonad (in the hermaphrodite, these structures are distal tip cells (DTCs), sheath cells, spermathecae, and uterus) and Z2 and Z3 give rise to the germ line. In later
25 larval stages the germ line contains a stem-cell population that contributes cells to

the meiotic pathway. Although germline nuclei reside in a syncytium at these stages, individual germline nuclei and their surrounding cytoplasm are typically referred to as a "germ cell". Development of the soma and germ line in both sexes, hermaphrodites and males, is coordinated by intercellular signaling. The self-fertile hermaphrodites are essentially modified females that produce sperm for a short time early in gametogenesis and then produce exclusively oocytes as adults. Males produce only sperm and can mate with hermaphrodites to produce cross progeny.

During post-embryonic development germ cells proliferate mitotically forming approximately 1000 nuclei in hermaphrodites and 500 in males. The *C. elegans* adult hermaphrodite gonad consists of two U-shaped gonad arms (FIG. 1). The two equivalent gonad arms of the adult hermaphrodite gonad have been described at an ultrastructural level (Hirsh *et al.*, 1976; Hall *et al.*, 1999; see FIG. 1). The distal portion of the gonad contains syncytial germline nuclei surrounded by incomplete membranes. The germ cells are connected to a core cytoplasm, also called the rachis. The stem-cell population is restricted to the distal-most part of the germ line; germ cells enter meiosis as they move proximally. In hermaphrodites, approximately the first 40 germ cells to enter meiotic prophase in each gonad arm differentiate as spermatocytes which complete meiosis to form approximately 160 sperm during the fourth larval stage of development. Upon progression to the adult stage, the germ cells differentiate as oocytes. Oocytes are surrounded by the proximal gonadal sheath cells (see FIG. 1).

The gonadal sheath cells are somatic cells that appear to play several roles important for the structure, integrity, and reproductive functions of the gonad (McCarter *et al.*, 1997; Rose *et al.*, 1997). The ten thin gonadal sheath cells can be subdivided into five pairs (1-5) with each pair having a distinct position along the proximal-distal axis of each gonad arm (FIG. 1). These elongated myoepithelial cells

lie between germ cells and the gonadal basal lamina (Hirsh *et al.*, 1976; Kimble and Hirsh, 1979; Strome, 1986; Hall *et al.*, 1999). The distal sheath cells (pair 1) have an unusual cellular structure with a flattened soma pressed into the gonad such that the cytoplasm is concentrated into a series of wedges that insert between the germ cells. Pair 1 distal sheath cells also extend finger-like filopodia between distal germ cells. Pair 2 ensheathes the loop region. The proximal sheath cells (pairs 3-5; see FIG. 1) contain thick and thin filaments and contract to drive ovulation (Strome, 1986; Myers *et al.*, 1996; McCarter *et al.*, 1997; Rose *et al.*, 1997; Hall *et al.*, 1999). The proximal sheath cells are positioned in an interdigitating pattern form gap junctions with one another, and are closely apposed to oocytes (Hall *et al.*, 1999). On their basal surfaces the proximal sheath cells attach to the gonadal basal lamina via hemi-adherens junctions which also serve to anchor the actin cytoskeleton and the contractile apparatus within the sheath cells. At their apical face, the proximal sheath cells often form gap junctions with oocytes. Yolk particles synthesized by the intestine (Kimble and Sharrock, 1983) gain access to oocytes for receptor-mediated endocytosis (Grant and Hirsh, 1999) by first moving through the sheath pores (Hall *et al.*, 1999). The most proximal sheath cells, pair 5, directly attach to the spermatheca. The spermatheca (1 per gonad arm) is a flexible accordion-like structure connected to the gonad arm distally and to the uterus proximally. The spermatheca expands greatly to accommodate oocytes, which are fertilized as they enter from the gonad arm during ovulation.

4.0 Female Sexual Maturation

The phrase "female sexual maturation" is defined herein to include, but is not limited to: meiotic maturation, completion of the meiotic divisions, oocyte production, oocyte or ovum maturation, and the events and processes of sheath cell

contraction and ovulation. In certain embodiments, "maturation" generally relates to the process by which an oocyte or an ova becomes competent for being fertilized. For example, female sexual maturation includes maturation of a female reproductive cell and maturation of an oocyte. In another example, as used herein,
5 female sexual maturation also includes a contraction of a sheath cell which is considered herein to be a female reproductive cell. Thus certain cells of the nematode reproductive system are referred to as female reproductive cells, even though they are not an oocyte or ovum per say.

5.0 Meiotic Maturation, Ovulation, and Completion of the Meiotic Divisions

10 Fully grown oocytes remain in the diakinesis stage of prophase I prior to undergoing meiotic maturation, ovulation, and fertilization. The nuclear envelope of the most proximal oocyte breaks down about 5 minutes prior to ovulation as it enters meiotic M-phase from prophase (Ward and Carrel, 1979; McCarter *et al.*, 1999). During maturation, the oocyte also undergoes a structural change termed
15 cortical rearrangement (McCarter *et al.*, 1999). These changes within the oocyte coincide with a reproducible sequence of somatic motor events mediated by the proximal sheath cells and the distal spermatheca resulting in ovulation. During ovulation the mature oocyte enters the spermatheca and is fertilized. The fertilized oocyte then passes into the uterus where both meiotic divisions are completed and
20 embryogenesis begins (Albertson, 1984; Albertson and Thomson, 1993; McCarter *et al.*, 1999). McCarter *et al.*, (1999) discloses that in the absence of sperm, the production of oocytes remains arrested in nematodes.

6.0 Assay to Identify the Stimulator of Female Sexual Maturation

The present inventors developed an *in vivo* assay for female sexual
25 maturation and used the assay to discover that the major sperm protein (MSP) is

the particular stimulator of female sexual maturation. The general procedures used are as follows. Large quantities of sperm (>10⁸) are purified using a modification of methods developed by Klass and Hirsh (1981). Synchronized cultures of *fog-2(q71)*, which are 50% male and 50% female, are used to purify adult males (Lewis and Flemming, 1995). Mutations in the *fog-2* gene block spermatogenesis in XX animals, transforming them into females, but have no effect on XO animals, which are fertile males (Schedl and Kimble, 1988). Males are separated from females, larvae, and embryos based on size by sieving through NITEX screens of various pore sizes. The populations of males isolated in this way are generally >99% pure. To isolate sperm, males are placed between two PLEXIGLASS plates and smashed in a vice grip (The Home Depot, Inc.). Intact sperm are then purified from the carcasses by filtration through NITEX filters (20 micron pore size) and washed in M9 phosphate buffer (Sulston and Hodgkin, 1988) using several rounds of low speed centrifugation and resuspension. Sperm-conditioned medium (SCM) are prepared by incubating purified sperm in M9 for different periods of time (e.g., 1-12 hours) and removing the sperm by centrifugation and filtration through a 0.2 micron filter. Microscopic analysis suggests that the sperm are not lysing during the incubation. Polyacrylamide gel electrophoresis (PAGE) also indicates that the sperm are not lysing during the incubation and also reveals that that an approximately 14 kDa protein is enriched in SCM. Referring to FIG. 2, a single protein band at approximately 14 kDa is apparent in the second lane. PAGE results for unfractionated SCM are displayed in lane 1.

Another aspect of the present invention provides an assay to screen SCM for maturation- and contraction-inducing activities, comprising microinjecting SCM into the uterus of reduced capacity sperm producing female nematodes (virgin *fog-2(q71)* females are used in certain preferred embodiments). Maturation and sheath

cell contraction are monitored by time-lapse video microscopy (see Rose *et al.* (1997) for a general description of time-lapse video microscopy of nematodes). Dramatic increases in oocyte maturation and sheath cell contraction rates are observed following injection of SCM. By contrast, no activity, or essentially no activity, is observed following injection of female extracts, 1-methyladenine, acetylcholine, oxytocin, or M9 buffer. The above described embodiment, provides a bioassay for sperm derived factors that promote oocyte maturation and sheath cell contraction. It is disclosed in the present invention that the activity is, at least in part, soluble and present in the sperm-conditioned media. This suggests that the factor is secreted by the sperm. Thus, it is a discovery of the present invention that while the sperm are sometime physically blocked by the valve, a soluble (diffusible) factor is secreted from the sperm to affect FSM and this factor penetrates the valve.

Still another aspect of the present invention provides compositions and methods for fractionation of SCM, and other nematode biological materials, for isolation of the FSM stimulatory factor. In one example, fractionation is performed using reversed phase high pressure liquid chromatography (HPLC) on Vydak C-4 and C-18 analytical columns. FIG. 3, a fraction marked by a + sign is the only active fraction recovered when the SCM is fractionated on the C-4 and C-18 columns, respectively (as determined using the *in vivo* FSM assay described herein). The biologically active fraction is analyzed using MALDI mass spectrometry peptide mapping and sequencing, identification techniques which are known in the art (FIG. 4). This result is verified by producing MSP-38 (GenBank Ac. # CAA93089) and MSP-142 (GenBank Ac. # CAB03037) in bacteria and purifying the respective isoforms using a commercially available 6-His tagging and protein product purification system. The inserts are cloned into the pQE-30 Type IV Kit available from Qiagen. The vector includes the 6-His tagging system and methods for this

cloning and use of the 6-His tagging system of identification and purification of expressed polypeptides are well known. The result is also confirmed using *C. elegans* MSP-77. Additional tests by MADLI-MS analysis demonstrate the *C. elegans* MSP in SCM is MSP-3 and MSP 142 (Miller *et al.*, 2001).

5 Microinjection of purified recombinant MSP mimics the biological response (FSM) seen using the active, fraction purified from SCM. Additionally, the recombinantly produced protein yields the same peptide map as the MSP from the active SCM fraction when cleaved with trypsin and analyzed using mass spectrometry. These results demonstrate that *C. elegans* sperm secrete, or
10 otherwise release, the major sperm protein, that the MSP signal is at least partially soluble in the extracellular fluid surrounds the sperm and in buffers, and that MSP dramatically increases FSM (including oocyte maturation, sheath cell contraction, and ovulation rates). The concept of MSP directed signal transduction of FSM including oocyte maturation, sheath cell contraction, and ovulation is diagrammed
15 in FIG. 5. Referring to FIG. 5, MSP serves as a simple molecule for communication between the sperm and non-mature or arrested oocytes and inactive female reproductive cells that sperm is present and ready to fertilize the egg.

Furthermore, the MSP is believed to be necessary and sufficient for stimulating that communication or signal transduction of FSM. While it is
20 contemplated and other factors may form a web-like network of upstream and/or downstream signal cascade, it is an advantage in certain embodiments herein that the role of MSP is quite uncomplicated. Thus, MSP provides an excellent target for anti-nematode agents. Also, due to the high sequence conservation between MSP polypeptides in all known nematodes, it is contemplated that nematodes will not
25 readily evolve resistance to compounds or agents that target MSP.

7.0 MSP is Known as a Structural Protein Localized Within the Sperm Cell

MSP is known in the art as a structural protein of the nematode sperm (Klass, M. R., *et al.*, 1981). Thus, it is a surprising discovery of the present invention that MSP is also a nematode female sexual maturation factor (stimulator, signal
5 transduction element, etc.).

It is widely accepted that motility of nematode sperm is not actin based, but rather is dependent upon MSP structure and function. Inside the sperm cell, dimeric MSP assembles at one end of a fibrous polymer of dimeric MSP and disassembles at the other end in a treadmill-like fashion which enables the sperm to
10 protrude and withdraw pseudopodia related to motility.

8.0 Sequence Conservation Among the Many MSP Sequences Described

There are likely more than sixty copies of the MSP gene in the *C. elegans* genome and it is believed that most of these MSP genes are transcribed. Referring to FIG. 6, twenty-seven MSP polypeptide sequences are provided corresponding to
15 polypeptides transcribed from apparently distinct MSP genes or polynucleotide sequences. Certain other nematodes apparently have fewer copies of MSP. For example, *A. suum* are believed to have two copies of an MSP gene both of which are believed to be transcribed into polypeptides.

Twenty-seven polypeptide sequences for FIG. 6 are aligned using Divide-and-
20 Conquer Multiple Sequence Alignment which is currently available over the world wide web (www) at the URL <http://bibiserv.techfak.uni-bielefeld.de/dca/>. The server is located at the Practical Computer Science and Bioinformatics research group which is run by Robert Giegerich. The physical location is: Robert Giegerich, AG
Praktische Informatik, Technische Fakultät, Universität Bielefeld, Postfach 10 01
25 31, D-33501 Bielefeld, Germany. The parameters used are Blossum 62 predefined

substitution matrix, free shift activated, approximate cut positions activated, recursion stop size L set to 20, window size W set to 0, and weight intensity lambda set to 0. The algorithm and method are disclosed in Stoye (1998).

Referring to FIG. 6, it is known that the N-terminal most methionine (from the ATG translation start site) is cleaved. It is expected that both forms (with and without the methionine) of MSP polypeptides are active in FSM; therefore, the methionine was included in FIG. 6. However, the references to the SEQ ID Numbers provided in FIG. 6 correspond to MSP polypeptide sequences of 126 amino acids and are without each N-terminal most methionine as shown in FIG. 6.

Again, referring to FIG. 6, the sequences of the numerous *C. elegans* MSP display a high degree of sequence homology. Very few sequence variations are observed. Residues that vary from the global (general) consensus within a column (determined visually) are marked in bold letter and underlined in FIG. 6. Because MSP polypeptide sequences, and even those of the most divergent known nematodes (see below), are so highly conserved; the preferred method for alignment is by visual inspection. For example, two or more MSP polypeptide sequences can be easily lined up next to one another on a computer screen or as written out on a paper and one moved against another until the majority of the bases match. Percent identity between any two sequences is calculated by counting the number of residues that do not match, dividing by the total number of residues in the total sequence being compared (or the shortest of the sequences being compared if one of the pair is shorter in length), multiplying by 100, and expressing the resulting value as a percent.

Using this approach, one of ordinary skill in the art can easily determine an alignment of a given MSP isolated from any member of the phylum Nematoda to another MSP, including to a *C. elegans* MSP, and in preferred embodiments the

MSP specified in SEQ ID NO:2. It is also preferred that the alignment selected is the one which produces the highest sequence identity as described above.

Thus, for example, the visual inspection method described above is used to align the sequences for the two known MSPs from *A. suum* (alpha and beta) with
5 MSP-142 of *C. elegans* (see FIG. 7). Again, very little sequence variation is observed between MSP polypeptides from these nematodes that are separated by hundreds of millions of years of evolutionary pressure. This indicates that the structure function relationship in MSP is tight and sequence variation likely results in a reduced fitness of the organism.

10 9.0 Biological Activities of MSP on FSM are Conserved in Phylum Nematoda

Ascaris suum is believed to be one of the most widely separated nematode species from *C. elegans* in terms of both evolution (hundreds of millions of years post divergence from the common organism) and in terms of distinctness of the MSP sequence including at the polypeptide level. For example, evaluation of the
15 biological activity of isolated MSP alpha and MSP beta from *A. suum* in the *in vivo* FSM assay described above serves as a model system for demonstrating that MSP sequences in general stimulate nematode FSM in all or nearly all member of the phylum Nematoda.

MSP, isoforms alpha from *A. suum* is isolated from *A. suum* nematodes or the
20 corresponding nucleotide is cloned and expressed in bacteria, for example. The specific sequence used is Accession Number P27439 in the NCBI database (maqsvppgdintqpsqkivfnapyddkhtyhikitnaggrrigwaikttnmrrlsvdppcgvldpkekvlmav scdtfnaatedlnndritiewtntpdgaakqfrrewfqgdgmvrknlpieynl) and is set forth in SEQ ID NO: 11, wherein SEQ ID NO:11 does not include the N-terminal methionine in

order to represent the polypeptide that is believed to be cleaved during cellular processing (see FIG. 7).

MSP, isoforms beta from *A. suum* is isolated from *A. suum* nematodes or the corresponding nucleotide is cloned and expressed in bacteria, for example. The
5 specific sequence used is Accession Number P27440
(maqsvppgdintqpgskivfnapyddkhtyhiktnaggrrigwaikttnmrrlgvdppcgvl dpkesvlmavs
cdtfnaatedlnndritiewtntpdgaakqfrrewfqgdgmvrknlpieynl) in the NCBI database
and is set forth in SEQ ID NO: 12, wherein SEQ ID NO:12 does not include the N-
terminal methionine in order to represent the polypeptide that is believed to be
10 cleaved during cellular processing.

Microinjection of MSP, isoform alpha or MSP, isoform beta into the sperm
defective (or reduced sperm capacity) female *C. elegans* as described above results in
a restoration of apparently normal FSM biological activity even though no sperm
are added. This provides support for using any combination of MSP polypeptide
15 including as expressed from MSP polynucleotides. Methods known in the art for
expressing polynucleotides that are preferred herein include, but are not limited to:
expression in bacteria, transient expression in nematodes, stable expression in
nematodes, and transgenic expression in nematodes. Transgenic expression in
nematodes includes gonadal specific expression and ectopic expression, such as in
20 transgenic expression in somatic cells of the nematode.

10.0 Experiments Demonstrating that MSP Stimulates FSM

The biological activities of MSP-77 and MSP 38 are studied using the *in vivo*
FSM assays described herein. Referring to FIG. 8, MSP isolated from male
nematodes, as well as MSP produced in bacteria stimulate oocyte maturation and
25 sheath cell contraction when introduced into female nematodes with reduced sperm

formation such as *fog-2* mutants (MSP-77 and MSP-38, in this figure, are isolated from the bacteria with a 6His tag. This general protein isolation technique is known in the art). Again, referring to FIG. 8, the top panel displays maturations per hour which is a biological measure of oocyte maturation. The center panel displays
5 contractions per minute which is a biological measure of sheath cell contraction. The bottom panel displays average displacement (in microns) which is another measure of sheath cell contraction. Measurements are made for buffer and the shown concentrations of MSP-77, MSP-38, and sperm MSP. The 6His marking denotes that the MSP includes a histidine tag (Qiagen) and was purified using this
10 system.

Similar experiments are performed with MSP-3 and MSP-142 (which are identified herein to be localized in sperm-conditioned medium), and MSP from *C. briggsae* and *C. remanei*. Also, experiments are performed with MSP from the distantly related *A. suum*. Each experiment demonstrates that MSP is essentially
15 interchangeable with regard to its biological activities in FSM.

11.0 The Carboxyl-Terminus of MSP Shows High Sequence Conservation

Residues 105 through 125 of MSPs derived from nine different genera of nematodes show a 100% sequence identity in these 19 consecutive amino acid residues (see FIG. 9). These nematodes represent free-living (*C. elegans*), animal
20 parasites (*Ascaris* and *Onchocerca*), and plant parasites (*Globodera*). (Specifying these genera as free-living, animal parasites, or plant parasites is not meant to limit the range that Nematodes inhabit the environment. In general many groups of nematodes have a diverse range)

12.0 Certain Domains Within MSP Differentially Stimulate FSM Activities

Another aspect of the present invention provides that certain domains within MSP differentially regulate certain FSM activities. For example, FIG. 10 demonstrates that residues 1-106 (SEQ ID NO:16) of MSP-77 (SEQ ID NO:9) preferentially stimulates oocyte maturation, while residues 106-126 (SEQ ID NO:17) of MSP-77 (SEQ ID NO:9) preferentially stimulates the rate of sheath cell contraction and displacement.

Thus, the biological activities of FSM, including oocyte maturation and sheath cell contraction, can be separated and the present invention discloses that different domains within MSP are capable of regulating those activities independently.

13.0 Description of MSP Sequence Fragments

Still another aspect of the present invention includes compositions and methods for identifying and using domains within MSP including for differential regulation of the biological activities of FSM. For example, specific sized segments of MSP polypeptide are systematically screened for the impact of that domain on FSM. This provides a fine resolution map of the MSP polypeptide with regard to FSM function that can be exploited to identify and manufacture highly specific anti-nematode agents, for example. This is also useful, for example, to identify FSM related domains of particularly high sequence conservation among members of Nematoda and/or to avoid areas that might include a short region that is similar to a gene or polypeptide in another organism, the targeting of which with anti-organism agents is not desired.

In certain embodiments antibodies (polyclonal and/or monoclonal) are generated against each fragment for use in labeling, identification, and an assay of

the present invention. In preferred embodiments, the antibodies are raised against those segments of MSP that influence FSM (either positively or negatively). In other embodiments, the antibodies are raised by injection of the MSP, or the FSM active domain of the MSP, into an animal (in certain embodiments, a non-human animal) using techniques known to one of skill in the art for producing antibodies. In other
5 embodiments, the antibodies and preferably monoclonal antibodies are produced in cell, such as hybridomas or by recombinant techniques that are known in the art. In certain embodiments, the antibodies are produced in humanized form. This can be accomplished in certain embodiments by injection of the immunogenic fragment of
10 MSP into a human.

14.0 Production of MSP Polypeptide Fragments

Segments of MSP polypeptide sequences are readily manufactured by chemical synthesis, for example by solid phase polypeptide manufacture. Alternatively, such segments can be cloned, synthesized in a heterologous cell
15 system, and isolated. Chemical synthesis is preferred for embodiments wherein the polypeptide is a polymer of 50 residues or fewer. Segments are polypeptides that are generally 10 amino acids or more in length (referring to consecutive amino acids in an MSP sequence in this section entitled Production of MSP Polypeptide Fragments). Although, in certain embodiments useful segments are contemplated
20 that include fewer than 10 consecutive MSP amino acids. It is believed herein that all or nearly all MSPs are essentially interchangeable with each other in regard to FSM activity. Although differences may be identified when examining the differential regulation of FSM activity.

Segments do not include the full length MSP polypeptide sequence. For
25 example, the 126 consecutive amino acids of SEQ ID NO:2 is not a segment.

Segments of a 126 amino acid MSP include 125 or fewer of consecutive amino acids of the MSP. It is preferred that the segment includes 120 or fewer, 110 or fewer, 106 or fewer, 105 or fewer, 100 or fewer, 95 or fewer, 90 or fewer, 85 or fewer, 80 or fewer, 75 or fewer, 70 or fewer, 65 or fewer, 60 or fewer, 55 or fewer, 50 or fewer, 45 or fewer, 40 or fewer, 30 or fewer, 25 or fewer, 20 or fewer, 19 or fewer, 18 or fewer, 17 or fewer, 16 or fewer, 15 or fewer, 14 or fewer, 13 or fewer, 12 or fewer, 11 or fewer, or 10 or fewer consecutive amino acids of an MSP. An MSP alignment variant means that amino acids may be substituted in any given MSP or MSP segment to make that position identical to the same position in another MSP molecule including from diverse groups of Nematoda. The segments may also be MSP alignment variant which may be referred to herein as MSP alignment variant segments. Functionally equivalent sequences and biological functional equivalents are also generally considered to be within the spirit and scope of the present invention and are described below.

Ranges of segments are also provided in certain embodiments of the present invention. For example, a 10 amino acid segment may be selected from any portion of the MSP polypeptide. An 11 amino acid segment may be selected from any portion of the MSP polypeptide. Segments of lengths including 9 amino acids (consecutive in an MSP), 10, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 106, 110, 115, 116, 117, 118, 119, 120, 121, 122, 123, and 124 are contemplated for certain embodiments.

These segments are generally screened for FSM activity or used in assays to identify MSP binding agents (e.g., agents that bind to specific domains) and to identify anti-nematode agents. Although many segments are described, they can be screened readily for FSM activity given the present disclosure and without undue

burden. A preferred method of screening is to utility high-throughput assays such as in microtitre plates or other multiple-sample format to rapidly examine a large number of segments. Such general assay are known in the art and are provided herein with regard to MSP and the embodiments of the present invention.

5 15.0 Description of Certain MSP Sequences

10 Numerous examples of MSP sequences (including polynucleotide and polypeptide) are known in the art and are useful for various embodiments of the present invention. Also, additional MSP sequences including from additional nematodes types and species can be readily determined, without burden, using conventional cloning techniques and the fact that many sequences are described. Thus, for example, primers can be designed to pull out additional polynucleotides from a pool, such as a gene or genomic library, including by PCR amplification. These polynucleotide can be expressed using standard cloning techniques and the MSP activity of the resulting polypeptide is measurable based upon the assays disclosed herein. Certain known and described MSPs are provided by way of example in Table 1 below. The MSPs are represented by Accession Numbers corresponding to files in the publicly available database maintained by the National Center for Biotechnology Information (the NCBI database). These files include polynucleotide sequences and polypeptide sequences of MSP. All information within
15
20 each file identified by Accession Number is hereby incorporated herein by reference.

The NCBI database is available on the world wide web at URL "http://www.ncbi.nlm.nih.gov/" and is physically located at: National Center for Biotechnology Information; National Library of Medicine; Building 38A, Room 8N805; Bethesda, MD 20894

Inventors: Greenstein and Miller
Attorney Docket: N-7088-MJM
Customer No. 23456

TABLE 1

Nematode Identifier	Nucleotide Accession Numbers	Protein Accession Numbers
<i>Mansonella ozzardi</i>	AJ404225	CAC20724
	AJ404224	CAC20723
	AJ404223	CAC20722
	AJ404222	CAC20721
	AJ404221	CAC20720
	AJ404220	CAC20719
	AJ404219	CAC20718
	AJ404218	CAC20717
	AJ404217	CAC20716
	AJ404216	CAC20715
	AJ404215	CAC20714
	AJ404214	CAC20713
	AJ404213	CAC20712
	AJ404212	CAC20711
	AJ404211	CAC20710
	AJ404210	CAC20709
	AJ404209	CAC20708
		CAC20742
<i>Onchocerca volvulus</i>	AJ404208	CAC20741
	AJ404207	CAC20740
	AJ404206	CAC20739
	AJ404205	CAC20738

Inventors: Greenstein and Miller
 Attorney Docket: N-7088-MJM
 Customer No. 23456

	AJ404204 J04663 J04662	B45528 A45528
<i>Ascaris suum</i>	X94249	A45944 P27439 P27440 AAB23264 CAA63933
<i>Ascaris lumbricoides</i>	M15680	AAA29375
<i>Globodera rostochiensis</i>	L24501 L24500 L24499	AAA29148 AAA29147 AAA29146
<i>Pratylenchus penetrans</i>		AAB02264 AAB02263 AAB02262 AAB02251 AAB02250 AAB02249
<i>Pratylenchus scribneri</i>		AAB02242 AAB02241 AAB02240 AAB02239

16.0 An Assay for Female Sexual Maturation

A method is provided of identifying an anti-nematode agent, by contacting a test compound to a nematode and monitoring a female sexual maturation of the nematode, wherein inhibition of the female sexual maturation indicates that the test compound includes the anti-nematode agent.

The *in vivo* bioassay is useful, for example, to identify sperm-related factors that promote oocyte maturation and gonadal sheath cell contraction. The assay is also useful, as another example, for identifying agents that inhibit nematode female sexual maturation.

In certain embodiments of the assay, mutant nematodes are utilized which have a reduced capacity for sperm production, or lack the capacity altogether. Such mutants are disclosed to have either low rates of oocyte maturation and sheath cell contraction activity or none at all. The present invention provides compositions and methods for using these mutants to screen for factors that stimulate or inhibit female sexual maturation. In similar embodiments, compositions and methods are provided that for using transgenic nematodes. These embodiments are described in detail below.

Test compounds include any compound in general. Preferred test compounds are soluble in aqueous solution and thus conducive to typical biological assay conditions. In certain embodiments, test compounds are selected from any chemical in a library, for example, as maintained by a pharmaceutical or other company. In certain embodiments, test compounds, include proteins, glycoproteins, polypeptides, glycopeptides, amino acids, nucleic acids of any variety, including DNA, RNA, peptide nucleic acid (PNA), carbohydrates, fatty acids, lipids, etc. In certain embodiments the test compound includes any biologically active molecule.

Test compounds are administered to nematodes by any desirable method for determining which of the test compounds has an inhibitory activity on nematode fertility, female sexual maturation, or other activity described herein. For example, the test compound can be microinjected, co-injected, incubated, or fed to the nematodes. The assay measures the ability of specific test compounds to inhibit MSP stimulation of oocyte maturation, gonadal sheath cell contraction, and ovulation using optical monitoring. In addition to ovulation, laying or releasing of oocytes or embryos from the organism can be monitored optically and used as an endpoint of test compound activity. The optical monitoring can be enhanced using labeling reagents, such as fluorescent, radioactive, or enzymatic labels. These labels can be attached using standard chemistry known in the art to the test compound, a sperm, a major sperm protein, the oocyte, a sheath cell, etc.

Methods for monitoring the FSM by video microscopy are disclosed herein. Other methods for monitoring the FSM can include by radiolabel assay, fluorescent assay, etc. (Miller *et al.*, Science 2001).

Inhibition of FSM generally refers to a reduction or termination in rate of an FSM event, in certain preferred embodiments. In other embodiments, inhibition means reduction in reproductive success, fecundity, etc. In other preferred embodiments, inhibition results in control of a population of nematodes including a free-living, parasitic, terrestrial, or an aquatic nematodes population.

17.0 Assay for Identifying Inhibitors of MSP Signaling

As described herein, an object of the present invention is to provide assays for screening test compounds to identify anti-nematode agents. In certain embodiments, the agents will be inhibitors of female nematode fertility. In certain embodiments, the agents will be inhibitors of an MSP signal transduction. Such

agents generally interfere with nematode fertility and are useful as agents for control of nematodes including free-living and animal and plant parasitic nematodes. Certain embodiments of screening assays for inhibitors of MSP signal transduction follow.

5 18.0 Mutant Nematode Strains

10 In general, MSP protein is administered to a mutant hermaphrodite or female nematode strain that does not produce sperm (*e.g.*, mutants in *fog-1*, *fog-2*, *fog-3*, *fem-1*, *fem-2*, *fem-3*, or *gld-1*(Fog). The MSP protein can be administered, for example, by microinjection into the uterus of the mutant hermaphrodite or female
15 nematode strain. Such strains are available from the *C. elegans* Genetic Center or natural isolates of gonochoristic species. In embodiments that use microinjection, the technique is carried out according to standard practice (*e.g.*, see LaMunyon and Ward, Genetics (1994) 138:689-692, incorporated herein by reference). Female nematode sexual maturation, such as oocyte maturation, sheath cell contraction, and ovulation are generally observed optically. Certain methods are described by McCarter *et al.*, *supra*.

19.0 Wild-Type Nematode Strains

20 In general, methods for screening test compounds for identifying factors that inhibit nematode fertility, and preferably that inhibit female sexual maturation, can be administered to wild-type nematodes and the effect of the test compounds is usually monitored optically. Typically, the test compounds can be co-injected, incubated, or fed to nematodes. Ordinarily the endpoint of the assay measures the ability of compounds to inhibit MSP stimulation of oocyte maturation, gonadal sheath cell contraction, and ovulation as described above.

20.0 Transgenic Nematode Strains

Nematode strains that ectopically express MSP (including in non-spermatogenic tissues) can be used in methods for screening test factors to identify anti-nematode agents. Transgenic nematodes expressing MSP are generated using standard methods (e.g., Methods in Cell Biology, ed. H. Epstein and D. Shakes, San Diego Academic Press, 452-482, incorporated by reference). Inhibitor compounds are, for example, incubated with or fed to the transgenic nematodes. The assay measures the ability of compounds to inhibit nematode fertility including, but not limited to: MSP stimulation of oocyte maturation, gonadal sheath cell contraction, and ovulation as described above. Typically, the inhibition of nematode fertility is measured optically. Test factors that inhibit MSP signal transduction will inhibit the nematode female sexual maturation. In addition, most transgenic strains produce different amounts of the transgenic factor, as is known in the art. Thus, nematodes that transgenically express varying amounts of MSP can be utilized to determine anti-nematode agent concentrations that are optimized for use on a particular nematode given that different nematodes express varying levels of MSP in the wild.

21.0 MSP Binding Assay

Compounds are screened for MSP binding affinity. Panels of candidate molecules are affixed to a matrix, for example microtitre wells, using standard methods. Labeled MSP protein, such as fluorescently, radioactively, or enzymatically linked MSP protein, is incubated with the compounds attached to the matrix, and then washed off (under conditions that remove unbound labeled MSP protein). Compounds which bind MSP are recognized by retention of the label (for example, optical recognition). Alternatively, MSP protein is affixed to a matrix, for

example microtitre wells, and incubated with labeled test compounds, such as fluorescently, radioactively, or enzymatically linked compounds, and then washed off, (under conditions that remove unbound labeled compound). Compounds which bind MSP are recognized by retention of the label.

5 In certain embodiments, compounds which bind MSP are then tested for the ability to block MSP signaling, for example by using bio-assays described herein. Compounds that inhibit or block nematode reproduction or fertility are anti-nematode agents. In certain embodiments, the anti-nematode agents are used to treat parasitic nematode infections in plants and animals by administering a
10 therapeutically effective amount of the anti-nematode agent to the plant or animal to inhibit, or in certain cases to virtually block, nematode reproduction in the infected plant or animal.

22.0 Regulation of MSP protein to protein interactions

15 Compounds are screened for the ability to regulate protein to protein interactions of MSP. For example, MSP is known to exist in monomeric and dimeric forms. Thus, test compounds are screened in biological assays to identify factors that prevent dimerization, multimerization (complexes with two or more MSP subunits), and for factors that prevent multimers from dissociating into monomers. The effect test compounds on multimer formation can be determined by incubating
20 the test compound with the MSP under multimer associating and dissociating conditions. These samples can be tested for biological activity in regard to nematode fertility as described herein. Multimer and monomer formation and dissociation can be monitored by techniques known in the art. For example, SDS versus native gel electrophoresis (polyacrylamide gel electrophoresis), electrospray mass
25 spectroscopy, and gel exclusion.

In certain embodiments, compounds which regulate multimerization of MSP (formation and dissociation of multimers/monomers) are then tested for the ability to block MSP signaling. For example by using biological assays measuring nematode female sexual maturation as described herein. Compounds that inhibit
5 nematode reproduction are anti-nematode agents. In certain embodiments, the anti-nematode agents are used to treat parasitic nematode infections in plants and animals by administering a therapeutically effective amount of the anti-nematode agent to the plant or animal to inhibit, or in certain cases to virtually block, nematode reproduction in the infected plant or animal.

10 23.0 Certain Nematode Varieties

Nematoda includes the roundworms and threadworms, and comprises a phylum of generally smooth-skinned, unsegmented worms with a long cylindrical body shape tapered at the ends; the phylum includes free-living and parasitic forms both aquatic and terrestrial (adapted from Academic Press Dictionary of Science
15 and Technology).

Table 2, below, provides a listing of the common name and scientific name of a multitude of nematode varieties. MSP genes and proteins can be derived from these or other nematode varieties and strains for use in conjunction with the present invention (*e.g.*, in a screening assay). Also, parasitic nematode infections of
20 these or other types of nematodes may be treated by anti-nematode agents described herein or identified as described herein. This list is not meant to be limited on the scope of the invention, but merely to be exemplary of types of nematode. Animal parasitic nematodes are also described in supplementary materials appended to this provisional application.

Table 2.

Common Name	Nematode Genus and species
African spiral nematode	<i>Helicotylenchus africanus</i>
Alfalfa root nematode	<i>Heterodera goettingiana</i>
Almond cyst nematode	<i>Heterodera amygdali</i>
Amaranth cyst nematode	<i>Cactodera amaranthi</i>
American dagger nematode	<i>Xiphinema americanum</i>
Amu-Darya nematode	<i>Heterodera oxiana</i>
Apple cyst nematode	<i>Globodera mali</i>
Apple root-knot nematode	<i>Meloidogyne mali</i>
Awl nematodes	<i>Dolichodorus</i> spp.
Banana meadow nematode	<i>Pratylenchus coffeae</i>
Banana nematode	<i>Pratylenchus musicola</i>
Banana root-lesion nematode	<i>Pratylenchus coffeae</i>
Banana spiral nematode	<i>Helicotylenchus multicinctus</i>
Banana-root nematode	<i>Radopholus similis</i>
Barley cyst nematode	<i>Heterodera hordecalis</i>
Barley root-knot nematode	<i>Meloidogyne nassi</i>
Beachgrass root-knot nematode	<i>Meloidogyne sasseri</i>
Beer nematode	<i>Panagrellus silusiae</i>
Beet nematode	<i>Heterodera schachtii</i>
Beet stem nematode	<i>Ditylenchus dipsaci</i>
Begonia leaf nematode	<i>Aphelenchoides fragariae</i>
Bentgrass nematode	<i>Anguina agrostis</i>
Bermudagrass cyst nematode	<i>Heterodera cardiolata</i>

Inventors: Greenstein and Miller
 Attorney Docket: N-7088-MJM
 Customer No. 23456

Birch cyst nematode	<i>Cactodera betulae</i>
Black currant nematode	<i>Aphelenchoides ritzemabosi</i>
Blueberry root-knot nematode	<i>Meloidogyne carolinensis</i>
Boxwood spiral nematode	<i>Rotylenchus buxophilus</i>
Brassica root eelworm	<i>Heterodera cruciferae</i>
Brassica root nematode	<i>Heterodera cruciferae</i>
Brazilian root-knot nematode	<i>Meloidogyne exigua</i>
Brazilian root-knot nematode	<i>Meloidogyne inornata</i>
British root-knot nematode	<i>Meloidogyne artiellia</i>
British spiral nematode	<i>Scutellonema brachyurum</i>
Buckwheat cyst nematode	<i>Heterodera graduni</i>
Bud and leaf nematodes	<i>Aphelenchoides</i> spp.
Bud and stem nematode	<i>Ditylenchus dipsaci</i>
Bulb and stem nematodes	<i>Ditylenchus</i> spp.
Bulb nematode	<i>Ditylenchus dipsaci</i>
Bulb or stem nematodes	<i>Ditylenchus</i> spp.
Burrowing nematode	<i>Radopholus similis</i>
Burrowing nematodes	<i>Radopholus</i> spp.
Cabbage cyst nematode	<i>Heterodera cruciferae</i>
Cabbage nematode	<i>Heterodera cruciferae</i>
Cabbage root nematode	<i>Heterodera cruciferae</i>
Cactus cyst nematode	<i>Cactodera cacti</i>
Cajanus cyst nematode	<i>Heterodera cajani</i>
California dagger nematode	<i>Xiphinema index</i>
California meadow nematode	<i>Pratylenchus neglectus</i>

Inventors: Greenstein and Miller
 Attorney Docket: N-7088-MJM
 Customer No. 23456

California root-lesion nematode	<i>Pratylenchus neglectus</i>
California sessile nematode	<i>Cacopaurus epacris</i>
Camel thorn cyst nematode	<i>Heterodera oxiana</i>
Camellia root-knot nematode	<i>Meloidogyne camelliae</i>
Canadian root-knot nematode	<i>Meloidogyne microtyla</i>
Carnation pin nematode	<i>Paratylenchus curvittatus</i>
Carnation pin nematode	<i>Paratylenchus dianthus</i>
Carolina spiral nematode	<i>Scutellonema brachyurum</i>
Carrot cyst nematode	<i>Heterodera carotae</i>
Carrot root nematode	<i>Heterodera carotae</i>
Cereal cyst nematode	<i>Heterodera avenae</i>
Cereal cyst nematode	<i>Heterodera latipons</i>
Cereal root nematode	<i>Heterodera avenae</i>
Cereal root-knot nematode	<i>Meloidogyne nassi</i>
Cereals root eelworm	<i>Heterodera major</i>
Cereals root nematode	<i>Heterodera avenae</i>
Chamber's dagger nematode	<i>Xiphinema chambersi</i>
Christie's spiral nematode	<i>Scutellonema christiei</i>
Christie's stubby root nematode	<i>Trichodorus christiei</i>
Chrysanthemum foliar nematode	<i>Aphelenchoides ritzemabosi</i>
Chrysanthemum leaf nematode	<i>Aphelenchoides ritzemabosi</i>
Chrysanthemum nematode	<i>Aphelenchoides ritzemabosi</i>
Citrus nematode	<i>Tylenchulus semipenetrans</i>
Citrus ring nematode	<i>Criconemoides citri</i>
Citrus root nematode	<i>Tylenchulus semipenetrans</i>

Inventors: Greenstein and Miller
 Attorney Docket: N-7088-MJM
 Customer No. 23456

Citrus root-knot nematode	<i>Meloidogyne indica</i>
Citrus spine nematode	<i>Criconema civellae</i>
Clover cyst nematode	<i>Heterodera trifolii</i>
Clover root nematode	<i>Heterodera trifolii</i>
Clover stem nematode	<i>Ditylenchus dipsaci</i>
Cobb's awl nematode	<i>Dolichodorus heterocephalous</i>
Cobb's lance nematode	<i>Hoplolaimus galeatus</i>
Cobb's meadow nematode	<i>Pratylenchus penetrans</i>
Cobb's ring nematode	<i>Criconemoides simile</i>
Cobb's root lesion nematode	<i>Pratylenchus penetrans</i>
Cobb's root-knot nematode	<i>Nacobbus batatiformis</i>
Cobb's spiral nematode	<i>Helicotylenchus multicinctus</i>
Cobb's stubby root nematode	<i>Trichodorus primitivus</i>
Coconut nematode	<i>Rhadinaphelenchus cocophilus</i>
Coconut palm nematode	<i>Rhadinaphelenchus cocophilus</i>
Cocopalm nematode	<i>Rhadinaphelenchus cocophilus</i>
Coffee meadow nematode	<i>Pratylenchus coffeae</i>
Coffee root-knot nematode	<i>Meloidogyne exigua</i>
Coffee root-lesion nematode	<i>Pratylenchus coffeae</i>
Columbia nematode	<i>Hoplolaimus columbus</i>
Columbia root-knot nematode	<i>Meloidogyne chitwoodi</i>
Corn cyst nematode	<i>Heterodera zeae</i>
Corn meadow nematode	<i>Pratylenchus zeae</i>
Corn root-lesion nematode	<i>Pratylenchus zeae</i>
Cotton root-knot nematode	<i>Meloidogyne incognita acrita</i>

Inventors: Greenstein and Miller
 Attorney Docket: N-7088-MJM
 Customer No. 23456

Cowpea cyst nematode	<i>Heterodera vigni</i>
Crown-headed lance nematode	<i>Hoplolaimus tylenchiformis</i>
Currant nematode	<i>Aphelenchoides ribes</i>
Cyperus cyst nematode	<i>Heterodera moths</i>
Cyst nematodes	<i>Globodera</i> spp.
Cyst nematodes	<i>Heterodera</i> spp.
Cyst-forming nematodes	<i>Heterodera</i> spp.
Cystoid body nematodes	<i>Meloidoderita</i> spp.
Cystoid nematodes	<i>Meloidodera</i> spp.
Dagger nematodes	<i>Xiphinema</i> spp.
De Man's meadow nematode	<i>Pratylenchus pratensis</i>
De Man's root-lesion nematode	<i>Pratylenchus pratensis</i>
Dock cyst nematode	<i>Heterodera rosii</i>
Douglas Fir nematode	<i>Nacobodera chitwoodi</i>
Ear-cockle nematode	<i>Anguina tritici</i>
Estonian cyst nematode	<i>Cactodera estonica</i>
Eucalypt cystoid nematode	<i>Cryphodera eucalypti</i>
European dagger nematode	<i>Xiphinema diversicaudatum</i>
False root-knot nematode of sugar beets	<i>Nacobbus batatiformis</i>
False root-knot nematode	<i>Nacobbus</i> spp.
Fern nematode	<i>Aphelenchoides fragariae</i>
Fern nematode	<i>Aphelenchoides olesistus</i>
Fescue leaf gall nematode	<i>Anguina graminis</i>
Ficus cyst nematode	<i>Heterodera fici</i>
Fig cyst nematode	<i>Heterodera fici</i>

Inventors: Greenstein and Miller
 Attorney Docket: N-7088-MJM
 Customer No. 23456

Fig pin nematode	<i>Paratylenchus hamatus</i>
Fig spine nematode	<i>Criconema decalineatum</i>
Foliar nematodes	<i>Aphelenchoides</i> spp.
Galeopsis cyst nematode	<i>Heterodera galeopsidis</i>
Galeopsis root nematode	<i>Heterodera galeopsidis</i>
Gall-forming nematodes	<i>Meloidogyne</i> spp.
Godfrey's meadow nematode	<i>Pratylenchus brachyurus</i>
Godfrey's root-lesion nematode	<i>Pratylenchus brachyurus</i>
Gold-plated nematode	<i>Globodera rostochiensis</i>
Golden nematode	<i>Globodera rostochiensis</i>
Golden nematode of potato	<i>Globodera rostochiensis</i>
Grass cyst nematode	<i>Punctodera punctata</i>
Grass root-gall nematode	<i>Subanguina radiculicola</i>
Grass sheath nematode	<i>Hemicycliophora similis</i>
Grass spiral nematode	<i>Helicotylenchus erythrinae</i>
Great root nematode	<i>Heterodera avenae</i>
Hairy-gall nematode	<i>Nacobbus batatiformis</i>
Heart-shaped cyst nematode	<i>Heterodera cardiolata</i>
Hop cyst nematode	<i>Heterodera humuli</i>
Hop nematode	<i>Heterodera humuli</i>
Hop root nematode	<i>Heterodera humuli</i>
Horsenettle cyst nematode	<i>Globodera tabacum virginiae</i>
Indian root-knot nematode	<i>Meloidogyne brevicauda</i>
Iris nematode	<i>Ditylenchus destructor</i>
Japanese cyst nematode	<i>Heterodera elachista</i>

Inventors: Greenstein and Miller
 Attorney Docket: N-7088-MJM
 Customer No. 23456

Javanese root-knot nematode	<i>Meloidogyne javanica</i>
Kansas cyst nematode	<i>Heterodera longicolla</i>
Kidney-shaped nematode	<i>Rotylenchulus reniformis</i>
Kikuyu grass nematode	<i>Meloidogyne kikuyuensis</i>
Knapweed nematode	<i>Mesoanguina picridis</i>
Knawel cyst nematode	<i>Heterodera scleranthii</i>
Knotweed cyst nematode	<i>Cactodera weissi</i>
Kona coffee root-knot nematode	<i>Meloidogyne konaensis</i>
Lance nematodes	<i>Hoplolaimus</i> spp.
Lesion nematodes	<i>Pratylenchus</i> spp.
Lespedeza cyst nematode	<i>Heterodera lespedezae</i>
Lucerne cyst nematode	<i>Heterodera medicaginis</i>
Maple root-knot nematode	<i>Meloidogyne ovalis</i>
Meadow nematodes	<i>Pratylenchus</i> spp.
Mediterranean cereal cyst nematode	<i>Heterodera latipons</i>
Milfoil cyst nematode	<i>Globodera millefolii</i>
Motha cyst nematode	<i>Heterodera mothi</i>
Mushroom nematode	<i>Aphelenchoides composticola</i>
Mushroom spawn nematode	<i>Ditylenchus myceliophagus</i>
Needle nematodes	<i>Longidorus</i> spp.
Nettle cyst nematode	<i>Heterodera urticae</i>
Nigerian dagger nematode	<i>Xiphinema nigeriense</i>
Northern root-knot nematode	<i>Meloidogyne hapla</i>
Nutgrass cyst nematode	<i>Heterodera cyperi</i>
Oak root-knot nematode	<i>Meloidogyne querciana</i>

Inventors: Greenstein and Miller
 Attorney Docket: N-7088-MJM
 Customer No. 23456

Oak sheathoid nematode	<i>Hemicriconemoides biformis</i>
Oat cyst nematode	<i>Heterodera avenae</i>
Oat cyst nematode	<i>Heterodera major</i>
Oat nematode	<i>Heterodera avenae</i>
Oat root nematode	<i>Heterodera avenae</i>
Onion stem nematode	<i>Ditylenchus dipsaci</i>
Osborne's cyst nematode	<i>Globodera tabacum solanacearum</i>
Pacific dagger nematode	<i>Xiphinema radicum</i>
Pea cyst nematode	<i>Heterodera goettingiana</i>
Pea root eelworm	<i>Heterodera goettingiana</i>
Pea root nematode	<i>Heterodera goettingiana</i>
Peanut root-knot nematode	<i>Meloidogyne arenaria</i>
Persian sessile nematode	<i>Cacopaurus pestis</i>
Phlox stem nematode	<i>Ditylenchus dipsaci</i>
Pigeon pea cyst nematode	<i>Heterodera cajani</i>
Pin nematodes	<i>Paratylenchus</i> spp.
Pine cystoid nematode	<i>Meloidodera floridensis</i>
Pine sheathoid nematode	<i>Hemicriconemoides floridensis</i>
Pine sting nematode	<i>Belonolaimus gracilis</i>
Pine wood nematode	<i>Bursaphelenchus xylophilus</i>
Potato cyst eelworm	<i>Globodera rostochiensis</i>
Potato cyst nematode	<i>Globodera pallida</i>
Potato cyst nematode	<i>Globodera rostochiensis</i>
Potato nematode	<i>Globodera rostochiensis</i>
Potato root eelworm	<i>Globodera rostochiensis</i>

Inventors: Greenstein and Miller
 Attorney Docket: N-7088-MJM
 Customer No. 23456

Potato root nematode	<i>Globodera rostochiensis</i>
Potato rot nematode	<i>Ditylenchus destructor</i>
Potato tuber eelworm	<i>Ditylenchus destructor</i>
Potato tuber nematode	<i>Ditylenchus destructor</i>
Pseudo root-knot nematode	<i>Hypsoperine graminis</i>
Ramie pin nematode	<i>Paratylenchus elachistus</i>
Red ring nematode	<i>Rhadinaphelenchus cocophilus</i>
Reniform nematode	<i>Rotylenchulus reniformis</i>
Reniform nematodes	<i>Rotylenchulus</i> spp.
Rice blind root nematodes	<i>Hirschmanniella</i> spp.
Rice cyst nematode	<i>Heterodera oryzae</i>
Rice nematode	<i>Aphelenchoides oryzae</i>
Rice root nematode	<i>Hirschmanniella oryzae</i>
Rice root-knot nematode	<i>Meloidogyne graminicola</i>
Rice stem nematode	<i>Ditylenchus angustus</i>
Rice stunt nematode	<i>Tylenchorhynchus martini</i>
Rice white-tip nematode	<i>Aphelenchoides besseyi</i>
Rice-root nematode	<i>Radopholus oryzae</i>
Ring nematodes	<i>Criconema</i> spp.
Ring nematodes	<i>Criconemoides</i> spp.
Root nematodes	<i>Heterodera</i> spp.
Root nematodes	<i>Hirschmanniella</i> spp.
Root-gall nematodes	<i>Meloidogyne</i> spp.
Root-knot nematodes	<i>Meloidogyne</i> spp.
Root-lesion nematodes	<i>Pratylenchus</i> spp.

Inventors: Greenstein and Miller
 Attorney Docket: N-7088-MJM
 Customer No. 23456

Round cystoid nematode	<i>Thecavermiculatus andinus</i>
Rubber cyst nematode	<i>Heterodera fici</i>
Rumex cyst nematode	<i>Heterodera rumicis</i>
Scribner's lesion nematode	<i>Pratylenchus scribneri</i>
Scribner's meadow nematode	<i>Pratylenchus scribneri</i>
Scribner's root-lesion nematode	<i>Pratylenchus scribneri</i>
Sedge cyst nematode	<i>Heterodera cyperi</i>
Seed gall nematode	<i>Afrina wevelli</i>
Seed gall nematodes	<i>Anguina</i> spp.
Seed-gall nematode	<i>Anguina tritici</i>
Seinhorst's stubby root nematode	<i>Trichodorus pachydermis</i>
Sessile nematodes	<i>Cacopaurus</i> spp.
Seville root-knot nematode	<i>Meloidogyne hispanica</i>
Sheath nematodes	<i>Hemicycliophora</i> spp.
Sheathoid nematodes	<i>Hemicriconemoides</i> spp.
Shoot gall nematodes	<i>Anguina</i> spp.
Smooth-headed lesion nematode	<i>Pratylenchus brachyurus</i>
Smooth-headed meadow nematode	<i>Pratylenchus leiocephalus</i>
Smooth-headed nematode	<i>Pratylenchus brachyurus</i>
Sorghum root-knot nematode	<i>Meloidogyne acronea</i>
Sour paste nematode	<i>Panagrellus redivivus</i>
South African pin nematode	<i>Paratylenchus curvittatus</i>
Southern root-knot nematode	<i>Meloidogyne incognita</i>
Sowthistle cyst nematode	<i>Heterodera sonchophila</i>
Soybean cyst nematode	<i>Heterodera glycines</i>

Spear nematodes	<i>Dorylaimus</i> spp.
Spine nematodes	<i>Criconema</i> spp.
Spiral nematodes	<i>Helicotylenchus</i> spp.
Spiral nematodes	<i>Rotylenchus</i> spp.
Spiral nematodes	<i>Scutellonema</i> spp.
Spring crimp nematode	<i>Aphelenchoides fragariae</i>
Spring dwarf nematode	<i>Aphelenchoides besseyi</i>
Steiner's spiral nematode	<i>Helicotylenchus dihystrera</i>
Stem gall nematode	<i>Pterotylenchus cecidogenus</i>
Stem nematode	<i>Ditylenchus dipsaci</i>
Sting nematode	<i>Belonolaimus gracilis</i>
Sting nematode	<i>Belonolaimus longicaudatus</i>
Sting nematodes	<i>Belonolaimus</i> spp.
Strawberry bud nematode	<i>Aphelenchoides besseyi</i>
Strawberry bud nematode	<i>Aphelenchoides fragariae</i>
Strawberry foliar nematode	<i>Aphelenchoides fragariae</i>
Strawberry nematode	<i>Aphelenchoides fragariae</i>
Stubby root nematode	<i>Trichodorus christiei</i>
Stubby root nematode	<i>Trichodorus kurumeensis</i>
Stubby root nematodes	<i>Paratrichodorus</i> spp.
Stubby root nematodes	<i>Trichodorus</i> spp.
Stunt nematode	<i>Tylenchorhynchus claytoni</i>
Stunt nematodes	<i>Tylenchorhynchus</i> spp.
Stylet nematodes	<i>Tylenchorhynchus</i> spp.
Sugar beet cyst nematode	<i>Heterodera schachtii</i>

Sugar beet nematode	<i>Heterodera schachtii</i>
Sugar cane cyst nematode	<i>Heterodera sacchari</i>
Sugar cane cyst nematode	<i>Heterodera schachtii</i>
Sugar cane stylet nematode	<i>Tylenchorhynchus martini</i>
Summer dwarf nematode	<i>Aphelenchoides fragariae</i>
Sycamore root-knot nematode	<i>Meloidogyne platani</i>
Tadzhik cyst nematode	<i>Heterodera tadshikistanica</i>
Tadzhik cystoid nematode	<i>Meloidodera tadshikistanica</i>
Tadzhik root-knot nematode	<i>Meloidogyne tadshikistanica</i>
Tarjan's sheath nematode	<i>Hemicycliophora parvana</i>
Tea root-knot nematode	<i>Meloidogyne brevicauda</i>
Teasel nematode	<i>Ditylenchus dipsaci</i>
Tesselate stylet nematode	<i>Tylenchorhynchus claytoni</i>
Thames' root-knot nematode	<i>Meloidogyne thamesi</i>
Thorne's cyst nematode	<i>Cactodera thornei</i>
Thorne's lance nematode	<i>Rotylenchus uniformis</i>
Thorne's meadow nematode	<i>Pratylenchus thornei</i>
Thorne's needle nematode	<i>Longidorus sylphus</i>
Thorne's root-lesion nematode	<i>Pratylenchus thornei</i>
Tobacco cyst nematode	<i>Globodera tabacum</i>
Tobacco stunt nematode	<i>Tylenchorhynchus claytoni</i>
Tulip root nematode	<i>Ditylenchus dipsaci</i>
Turf spiral nematode	<i>Rotylenchus christiei</i>
Turkmen cyst nematode	<i>Heterodera turcomanica</i>
Ustinov cyst nematode	<i>Heterodera ustinovi</i>

Valentine cyst nematode	<i>Heterodera cardiolata</i>
Vinegar eels	<i>Turbatrix aceti</i>
Vinegar nematode	<i>Turbatrix aceti</i>
Walnut meadow nematode	<i>Pratylenchus vulnus</i>
Walnut root-lesion nematode	<i>Pratylenchus vulnus</i>
Walnut sessile nematode	<i>Cacopaurus pestis</i>
Wesson's sheathoid nematode	<i>Hemicriconemoides wessoni</i>
West African spiral nematode	<i>Scutellonema blaberum</i>
Wheat cyst nematode	<i>Heterodera latipons</i>
Wheat gall nematode	<i>Anguina tritici</i>
White-tip nematode	<i>Aphelenchoides besseyi</i>
Willow cyst nematode	<i>Heterodera salixophila</i>
Yam nematode	<i>Scutellonema bradys</i>
Yarrow cyst nematode	<i>Globodera achilleae</i>
Zimmerman's spiral nematode	<i>Helicotylenchus erythrinae</i>
Zoysia spine nematode	<i>Criconema spinalineatum</i>

24.0 Treating a parasitic nematode infection

In certain embodiments of the present invention, a parasitic nematode infection is treated in an infected organism (including plants and animals). A preferred method of treating a parasitic nematode infection is to inhibit nematode fertility or reproduction in the infected animal. In general, this is done by administering a therapeutically effective amount of an anti-nematode agent as described herein which disrupts a biological activity of MSP related to female

nematode sexual maturation. The anti-nematode agent can be identified as described in the present invention.

5 In certain embodiments, a method of inhibiting a reproduction of a nematode is provided, comprising inhibiting a signal transduction of a major sperm protein of the nematode, wherein the signal transduction stimulates a female sexual maturation.

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

In certain embodiments, MSP signal transduction is inhibited by a method comprising administering an MSP-specific antibody (Several MSP-specific antibodies have been described in the art. For example, see, Klass, M. R., and Hirsh, D. 1981. Sperm isolation and biochemical analysis of the major sperm protein from *Caenorhabditis elegans*. Dev. Biol. 84, 299-312, incorporated herein by reference). (In certain examples, the MSP-specific antibody can be administered by solublizing the antibody in the nematode's environment or by microinjection into the uterus of the nematode. Without being bound to mechanism or theory, these antibodies bind MSP *in vivo* and inhibit MSP-mediated signaling. Experiments are contemplated wherein hermaphrodite oocyte maturation and sheath cell contraction rates will be compared to hermaphrodites injected with an antibody which does not bind MSP (a suitable antibody for a negative control experiment).

20 In certain embodiments, MSP signal transduction is inhibited by administering a vaccine (especially in the case of treating an animal) wherein the vaccine is developed to an MSP protein or fragment thereof. Any variety of MSP protein, or fragment thereof, can be used in vaccine development. Many nematode types are described herein and the MSP or MSPs isolated therefrom may be used. Vaccine development protocols are well known.

25 In certain embodiments the identified anti-nematode agent is applied to crops including by spraying a field to distribute the agent. The agent may gain access

directly to the nematode or exist in the soil, water, food or general environment of the nematode. The agent may also be transferred to the nematode from a plant or animal.

25.0 Pharmaceutical formulations

5 In certain embodiments, the preferred method of administering a biologically active molecule (such as MSP or an anti-nematode agent) is in combination with an excipient (a pharmaceutically acceptable carrier). The combination of at least one pharmaceutically acceptable carrier and at least one biologically active molecule is referred to herein as a pharmaceutical formulation.

10 The particular excipient is not believed to be critical as long as it is compatible with the biological activity of the biologically active molecule and compatible with administration to the subject, especially plants, animals, human, and other mammals. The choice of excipient depends on the nature of the treatment being administered and the biologically active molecule. The pharmaceutical formulation can be applied to the surface of the organism being treated for a parasitic nematode infection or injected into the local tissue either in one application or multiple applications. The pharmaceutical formulation can be combined with additional inert or carrier ingredients and used as a topical salve. The pharmaceutical formulation may also be aerosolized and administered through
20 the lungs.

In certain preferred embodiments, a pharmaceutical formulation is provided along with a method of applying a metered amount of the formulation. For example, if a syringe may include unit markings on the barrel of the syringe. Typically a syringe will also include a needle and a plunger to form a device effective for
25 administration by injection. The particular choice of pharmaceutically acceptable

carrier can be made by one with skill in the art, such as a treating physician, veterinarian, farmer, or extension service personnel.

As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic
5 and absorption delaying agents and the like that are compatible with the biologically active molecule(s) and with administration to the organism being treated.

In certain embodiments, the pharmaceutical formulations of the present invention are advantageously administered either as liquid solutions or suspensions. Solid forms may be solubilized or suspended in liquid prior to application or injection. These preparations also may be emulsified. In certain
10 embodiments, a typical composition comprises about 50 mg or up to about 100 mg of human serum albumin per milliliter of phosphate buffered saline. Other pharmaceutically acceptable carriers include aqueous solutions, salts, preservatives, buffers and the like. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oil, and organic esters, such as theyloleate. Aqueous carriers include water, alcoholic/aqueous solutions, and saline solutions including sodium chloride, Ringer's dextrose, etc. Preservatives include antimicrobial agents, anti-oxidants, chelating agents and inert gases. The pH and
15 20 exact concentration of the various components in the pharmaceutical formulation are adjusted according to well known parameters using well known buffering and dilution agents.

26.0 Production of Recombinant MSP Containing Vectors

Sub A57
25 Recombinant MSP bacterial strains were produced by cloning MSP-142 and MSP-38 into the pQe-30 6-His vector from Qiagen. Primers specific for MSP were

sub A's
cont.

made that contained a 5' BamHI site (5' primer) or a 5' HindIII (3' primer) followed by the respective MSP-coding sequences. MSP-38 and MSP-142 were amplified by PCR, cut with BamHI and HindIII, and ligated into the pQe-30 vector (FIG. 6) which was also cut with BamHI and HindIII. This strategy generated a vector
5 containing an IPTG-inducible promoter followed by an initiator methionine, an N-terminal 6-His tag, and the respective MSP-38 or MSP-142 coding sequences. This construct was then transformed into M15(pREP4) bacterial cells and vector-containing colonies were selected with LB medium containing Ampicillin and Kanamycin. MSP-containing colonies were grown overnight and then MSP expression was induced for 4 hours with 1 mM IPTG. Induced bacteria were pelleted, lysed, and purified using a NiNTA agarose column, which binds the 6-His tag. 6-His purification is known in the art.

27.0 Biological Functional Equivalents of Polynucleotides and Polypeptides

As is known to one with skill in the art, the biological function or activity of a gene product may not correspond directly to an absolute polynucleotide or polypeptide sequence of the gene product. Therefore, the inventor specifically contemplates that alterations to sequences provided herein may be made or used wherein the altered sequences, or methods of use thereof, are equivalent to sequences, or methods of use thereof, and are within the spirit and scope of the
20 present invention. These equivalent sequences are referred to as biologically functional equivalents, or simply as functional equivalents. Functional equivalents can include, but are not limited to: conservatively modified variants, degeneracy of the nucleic acid code, polymorphisms, certain insertions and deletions, and certain length variants. Methods for altering sequence residues and testing the altered

sequences for function or activity are known in the art or described herein. These alterations may be natural or made by the "hand of man".

At the nucleotide level, different codons can encode the same amino acid. In other words, the genetic code is degenerate (Alberts *et al.*, Molecular Biology of the Cell, (1989) 2nd Edition, Garland Publishing, Inc., and incorporated herein by reference). The terms "wobble" and "nucleic acid degeneracy" are used herein to refer to codons that encode the same amino acid, such as the six codons for arginine or serine. Preferred human codons are provided in materials appended to this application. Codon preferences for other organisms also are well known to those of skill in the art (Wada *et al.*, 1990, *supra*). Thus, one with skill in the art knows that two different polynucleotides can encode identical polypeptide sequences due to codon wobble.

It is understood in the art that amino acid and nucleic acid sequences may include additional residues, such as additional N-terminal or C-terminal amino acids or 5' or 3' sequences, and yet still be essentially as set forth in one of the sequences disclosed herein; so long as the sequence meets the criteria set forth herein, including the maintenance of at least one biological protein activity where protein expression is concerned (MSP activity should include at least one type of stimulation of female nematode sexual maturation). The addition of terminal sequences particularly applies to nucleic acid sequences that may, for example, include various non-coding sequences flanking either of the 5' or 3' portions of the coding region or may include various internal sequences, *i.e.*, introns, which are known to occur within genes between coding regions (Alberts *et al.*, *supra*, incorporated herein by reference). Thus; about 1, 2, 3, 4, 5, 6, 7, or more than 7 amino acids could be added to a polypeptide and the polypeptide may still retain at least one biological activity. Or; about 1, 2, 3, 4, 5, 6, 7, or more than 7 nucleotides

could be added to a polynucleotide and expression products of the polynucleotide may still retain at least one biological activity.

5 It also is understood in the art that amino acid and nucleic acid residues may be removed from the N-terminal or C-terminal ends of polypeptide or 5' or 3' ends of polynucleotide sequences, and yet still be essentially as set forth in one of the sequences disclosed herein; so long as the sequence meets the criteria set forth herein, including the maintenance of at least one biological protein activity of where protein expression is concerned (in particular stimulating a female nematode sexual maturation with regard to MSP). The removal of terminal sequences particularly applies to nucleic acid sequences that may, for example, include various non-coding sequences flanking either of the 5' or 3' portions of the coding region or may include various internal sequences, *i.e.*, introns, which are known to occur within genes between coding regions (Alberts *et al.*, *supra*, incorporated herein by reference). Thus; about 1, 2, 3, 4, 5, 6, 7, or more than 7 amino acids could be removed from a polypeptide and the polypeptide may still retain at least one biological activity. Or, about 1, 2, 3, 4, 5, 6, 7, or more than 7 nucleotides could be removed from a polynucleotide and expression products of the polynucleotide may still retain at least one biological activity.

10
15
20
25 If desired, it is possible using techniques known to one with skill in the art, to include an intron in a recombinant polynucleotide sequence. For example, a bovine growth hormone (bGH) intron including splice sites may be added. In certain instances, the addition of an intron to a recombinant polynucleotide has been observed to increase expression of the encoded expression product in eukaryotic cells. It is understood that the addition of an intron may create a functionally equivalent sequence.

It is further understood in the art that insertions and deletions may be made within the amino acid and nucleic acid sequence, and yet still be essentially as set forth in one of the sequences disclosed herein; so long as the sequence meets the criteria set forth herein, including the maintenance of biological protein activity
5 where protein expression is concerned (for MSP this should be a stimulation of at least one female nematode sexual maturation). It is preferred that the reading frame of a polynucleotide sequence be maintained, as is known in the art (Alberts *et al.*, supra, incorporated herein by reference).

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
1000
1005
1010
1015
1020
1025
1030
1035
1040
1045
1050
1055
1060
1065
1070
1075
1080
1085
1090
1095
1100
1105
1110
1115
1120
1125
1130
1135
1140
1145
1150
1155
1160
1165
1170
1175
1180
1185
1190
1195
1200
1205
1210
1215
1220
1225
1230
1235
1240
1245
1250
1255
1260
1265
1270
1275
1280
1285
1290
1295
1300
1305
1310
1315
1320
1325
1330
1335
1340
1345
1350
1355
1360
1365
1370
1375
1380
1385
1390
1395
1400
1405
1410
1415
1420
1425
1430
1435
1440
1445
1450
1455
1460
1465
1470
1475
1480
1485
1490
1495
1500
1505
1510
1515
1520
1525
1530
1535
1540
1545
1550
1555
1560
1565
1570
1575
1580
1585
1590
1595
1600
1605
1610
1615
1620
1625
1630
1635
1640
1645
1650
1655
1660
1665
1670
1675
1680
1685
1690
1695
1700
1705
1710
1715
1720
1725
1730
1735
1740
1745
1750
1755
1760
1765
1770
1775
1780
1785
1790
1795
1800
1805
1810
1815
1820
1825
1830
1835
1840
1845
1850
1855
1860
1865
1870
1875
1880
1885
1890
1895
1900
1905
1910
1915
1920
1925
1930
1935
1940
1945
1950
1955
1960
1965
1970
1975
1980
1985
1990
1995
2000
2005
2010
2015
2020
2025
2030
2035
2040
2045
2050
2055
2060
2065
2070
2075
2080
2085
2090
2095
2100
2105
2110
2115
2120
2125
2130
2135
2140
2145
2150
2155
2160
2165
2170
2175
2180
2185
2190
2195
2200
2205
2210
2215
2220
2225
2230
2235
2240
2245
2250
2255
2260
2265
2270
2275
2280
2285
2290
2295
2300
2305
2310
2315
2320
2325
2330
2335
2340
2345
2350
2355
2360
2365
2370
2375
2380
2385
2390
2395
2400
2405
2410
2415
2420
2425
2430
2435
2440
2445
2450
2455
2460
2465
2470
2475
2480
2485
2490
2495
2500
2505
2510
2515
2520
2525
2530
2535
2540
2545
2550
2555
2560
2565
2570
2575
2580
2585
2590
2595
2600
2605
2610
2615
2620
2625
2630
2635
2640
2645
2650
2655
2660
2665
2670
2675
2680
2685
2690
2695
2700
2705
2710
2715
2720
2725
2730
2735
2740
2745
2750
2755
2760
2765
2770
2775
2780
2785
2790
2795
2800
2805
2810
2815
2820
2825
2830
2835
2840
2845
2850
2855
2860
2865
2870
2875
2880
2885
2890
2895
2900
2905
2910
2915
2920
2925
2930
2935
2940
2945
2950
2955
2960
2965
2970
2975
2980
2985
2990
2995
3000
3005
3010
3015
3020
3025
3030
3035
3040
3045
3050
3055
3060
3065
3070
3075
3080
3085
3090
3095
3100
3105
3110
3115
3120
3125
3130
3135
3140
3145
3150
3155
3160
3165
3170
3175
3180
3185
3190
3195
3200
3205
3210
3215
3220
3225
3230
3235
3240
3245
3250
3255
3260
3265
3270
3275
3280
3285
3290
3295
3300
3305
3310
3315
3320
3325
3330
3335
3340
3345
3350
3355
3360
3365
3370
3375
3380
3385
3390
3395
3400
3405
3410
3415
3420
3425
3430
3435
3440
3445
3450
3455
3460
3465
3470
3475
3480
3485
3490
3495
3500
3505
3510
3515
3520
3525
3530
3535
3540
3545
3550
3555
3560
3565
3570
3575
3580
3585
3590
3595
3600
3605
3610
3615
3620
3625
3630
3635
3640
3645
3650
3655
3660
3665
3670
3675
3680
3685
3690
3695
3700
3705
3710
3715
3720
3725
3730
3735
3740
3745
3750
3755
3760
3765
3770
3775
3780
3785
3790
3795
3800
3805
3810
3815
3820
3825
3830
3835
3840
3845
3850
3855
3860
3865
3870
3875
3880
3885
3890
3895
3900
3905
3910
3915
3920
3925
3930
3935
3940
3945
3950
3955
3960
3965
3970
3975
3980
3985
3990
3995
4000
4005
4010
4015
4020
4025
4030
4035
4040
4045
4050
4055
4060
4065
4070
4075
4080
4085
4090
4095
4100
4105
4110
4115
4120
4125
4130
4135
4140
4145
4150
4155
4160
4165
4170
4175
4180
4185
4190
4195
4200
4205
4210
4215
4220
4225
4230
4235
4240
4245
4250
4255
4260
4265
4270
4275
4280
4285
4290
4295
4300
4305
4310
4315
4320
4325
4330
4335
4340
4345
4350
4355
4360
4365
4370
4375
4380
4385
4390
4395
4400
4405
4410
4415
4420
4425
4430
4435
4440
4445
4450
4455
4460
4465
4470
4475
4480
4485
4490
4495
4500
4505
4510
4515
4520
4525
4530
4535
4540
4545
4550
4555
4560
4565
4570
4575
4580
4585
4590
4595
4600
4605
4610
4615
4620
4625
4630
4635
4640
4645
4650
4655
4660
4665
4670
4675
4680
4685
4690
4695
4700
4705
4710
4715
4720
4725
4730
4735
4740
4745
4750
4755
4760
4765
4770
4775
4780
4785
4790
4795
4800
4805
4810
4815
4820
4825
4830
4835
4840
4845
4850
4855
4860
4865
4870
4875
4880
4885
4890
4895
4900
4905
4910
4915
4920
4925
4930
4935
4940
4945
4950
4955
4960
4965
4970
4975
4980
4985
4990
4995
5000
5005
5010
5015
5020
5025
5030
5035
5040
5045
5050
5055
5060
5065
5070
5075
5080
5085
5090
5095
5100
5105
5110
5115
5120
5125
5130
5135
5140
5145
5150
5155
5160
5165
5170
5175
5180
5185
5190
5195
5200
5205
5210
5215
5220
5225
5230
5235
5240
5245
5250
5255
5260
5265
5270
5275
5280
5285
5290
5295
5300
5305
5310
5315
5320
5325
5330
5335
5340
5345
5350
5355
5360
5365
5370
5375
5380
5385
5390
5395
5400
5405
5410
5415
5420
5425
5430
5435
5440
5445
5450
5455
5460
5465
5470
5475
5480
5485
5490
5495
5500
5505
5510
5515
5520
5525
5530
5535
5540
5545
5550
5555
5560
5565
5570
5575
5580
5585
5590
5595
5600
5605
5610
5615
5620
5625
5630
5635
5640
5645
5650
5655
5660
5665
5670
5675
5680
5685
5690
5695
5700
5705
5710
5715
5720
5725
5730
5735
5740
5745
5750
5755
5760
5765
5770
5775
5780
5785
5790
5795
5800
5805
5810
5815
5820
5825
5830
5835
5840
5845
5850
5855
5860
5865
5870
5875
5880
5885
5890
5895
5900
5905
5910
5915
5920
5925
5930
5935
5940
5945
5950
5955
5960
5965
5970
5975
5980
5985
5990
5995
6000
6005
6010
6015
6020
6025
6030
6035
6040
6045
6050
6055
6060
6065
6070
6075
6080
6085
6090
6095
6100
6105
6110
6115
6120
6125
6130
6135
6140
6145
6150
6155
6160
6165
6170
6175
6180
6185
6190
6195
6200
6205
6210
6215
6220
6225
6230
6235
6240
6245
6250
6255
6260
6265
6270
6275
6280
6285
6290
6295
6300
6305
6310
6315
6320
6325
6330
6335
6340
6345
6350
6355
6360
6365
6370
6375
6380
6385
6390
6395
6400
6405
6410
6415
6420
6425
6430
6435
6440
6445
6450
6455
6460
6465
6470
6475
6480
6485
6490
6495
6500
6505
6510
6515
6520
6525
6530
6535
6540
6545
6550
6555
6560
6565
6570
6575
6580
6585
6590
6595
6600
6605
6610
6615
6620
6625
6630
6635
6640
6645
6650
6655
6660
6665
6670
6675
6680
6685
6690
6695
6700
6705
6710
6715
6720
6725
6730
6735
6740
6745
6750
6755
6760
6765
6770
6775
6780
6785
6790
6795
6800
6805
6810
6815
6820
6825
6830
6835
6840
6845
6850
6855
6860
6865
6870
6875
6880
6885
6890
6895
6900
6905
6910
6915
6920
6925
6930
6935
6940
6945
6950
6955
6960
6965
6970
6975
6980
6985
6990
6995
7000
7005
7010
7015
7020
7025
7030
7035
7040
7045
7050
7055
7060
7065
7070
7075
7080
7085
7090
7095
7100
7105
7110
7115
7120
7125
7130
7135
7140
7145
7150
7155
7160
7165
7170
7175
7180
7185
7190
7195
7200
7205
7210
7215
7220
7225
7230
7235
7240
7245
7250
7255
7260
7265
7270
7275
7280
7285
7290
7295
7300
7305
7310
7315
7320
7325
7330
7335
7340
7345
7350
7355
7360
7365
7370
7375
7380
7385
7390
7395
7400
7405
7410
7415
7420
7425
7430
7435
7440
7445
7450
7455
7460
7465
7470
7475
7480
7485
7490
7495
7500
7505
7510
7515
7520
7525
7530
7535
7540
7545
7550
7555
7560
7565
7570
7575
7580
7585
7590
7595
7600
7605
7610
7615
7620
7625
7630
7635
7640
7645
7650
7655
7660
7665
7670
7675
7680
7685
7690
7695
7700
7705
7710
7715
7720
7725
7730
7735
7740
7745
7750
7755
7760
7765
7770
7775
7780
7785
7790
7795
7800
7805
7810
7815
7820
7825
7830
7835
7840
7845
7850
7855
7860
7865
7870
7875
7880
7885
7890
7895
7900
7905
7910
7915
7920
7925
7930
7935
7940
7945
7950
7955
7960
7965
7970
7975
7980
7985
7990
7995
8000
8005
8010
8015
8020
8025
8030
8035
8040
8045
8050
8055
8060
8065
8070
8075
8080
8085
8090
8095
8100
8105
8110
8115
8120
8125
8130
8135
8140
8145
8150
8155
8160
8165
8170
8175
8180
8185
8190
8195
8200
8205
8210
8215
8220
8225
8230
8235
8240
8245
8250
8255
8260
8265
8270
8275
8280
8285
8290
8295
8300
8305
8310
8315
8320
8325
8330
8335
8340
8345
8350
8355
8360
8365
8370
8375
8380
8385
8390
8395
8400
8405
8410
8415
8420
8425
8430
8435
8440
8445
8450
8455
8460
8465
8470
8475
8480
8485
8490
8495
8500
8505
8510
8515
8520
8525
8530
8535
8540
8545
8550
8555
8560
8565
8570
8575
8580
8585
8590
8595
8600
8605
8610
8615
8620
8625
8630
8635
8640
8645
8650
8655
8660
8665
8670
8675
8680
8685
8690
8695
8700
8705
8710
8715
8720
8725
8730
8735
8740
8745
8750
8755
8760
8765
8770
8775
8780
8785
8790
8795
8800
8805
8810
8815
8820
8825
8830
8835
8840
8845
8850
8855
8860
8865
8870
8875
8880
8885
8890
8895
8900
8905
8910
8915
8920
8925
8930
8935
8940
8945
8950
8955
8960
8965
8970
8975
8980
8985
8990
8995
9000
9005
9010
9015
9020
9025
9030
9035
9040
9045
9050
9055
9060
9065
9070
9075
9080
9085
9090
9095
9100
9105
9110
9115
9120
9125
9130
9135
9140
9145
9150
9155
9160
9165
9170
9175
9180
9185
9190
9195
9200
9205
9210
9215
9220
9225
9230
9235
9240
9245
9250
9255
9260
9265
9270
9275
9280
9285
9290
9295
9300
9305
9310
9315
9320
9325
9330
9335
9340
9345
9350
9355
9360
9365
9370
9375
9380
9385
9390
9395
9400
9405
9410
9415
9420
9425
9430
9435
9440
9445
9450
9455
9460
9465
9470
9475
9480
9485
9490
9495
9500
9505
9510
9515
9520
9525
9530
9535
9540
9545
9550
9555
9560
9565
9570
9575
9580
9585
9590
9595
9600
9605
9610
9615
9620
9625
9630
9635
9640
9645
9650
9655
9660
9665
9670
9675
9680
9685
9690
9695
9700
9705
9710
9715
9720
9725
9730
9735
9740
9745
9750
9755
9760
9765
9770
9775
9780
9785
9790
9795
9800
9805
9810
9815
9820
9825
9830
9835
9840
9845
9850
9855
9860
9865
9870
9875
9880
9885
9890
9895
9900
9905
9910
9915
9920
9925
9930
9935
9940
9945
9950
9955
9960
9965
9970
9975
9980
9985
9990
9995
10000
10005
10010
10015
10020
10025
10030
10035
10040
10045
10050
10055
10060
10065
10070
10075
10080
10085
10090
10095
10100
10105
10110
10115
10120
10125
10130
10135
10140
10145
10150
10155
10160
10165
10170
10175
10180
10185
10190
10195
10200
10205
10210
10215
10220
10225
10230
10235
10240
10245
1025

interactions are well known in the art including techniques such as SDS-PAGE and Western analysis.

10 The nucleic acid segments of the present invention, regardless of the length of the coding sequence itself, may be combined with other DNA sequences (one or
5 more of each), such as promoters, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, internal ribosome entry sites, introns, other coding segments, membrane transport sequences, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol. Therefore, the terms "MSP gene" may also comprise any combination of associated control sequences. Furthermore, those skilled in the art of mutagenesis will appreciate that other analogs, as yet undisclosed or undiscovered, may be used to construct MSP analogs (mutants, variants, etc). Additional meaning
15 of biological functional equivalents, similarity, percent similarity, equivalents, substantially identical sequences, essentially the same, and essentially similar sequences and activities are described in U.S. Patent 5,922,688 to Hung *et al.*, incorporated herein by reference.

20 Naturally, the present invention also encompasses peptides and polypeptides (or the nucleic acid sequences that encode such peptides and polypeptides) that contain conservatively modified variants of sequences of interest, for example, a MSP sequence. One with skill in the art is able to determine conservative sequence modifications. In the case of a polypeptide, amino acid substitutions, such as those which might be employed in modifying a peptide, such as MSP, are generally based
25 on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like.

An analysis of the size, shape and type of the amino acid side-chain substituents reveals that arginine, lysine and histidine are all positively charged residues; that alanine, glycine and serine are all a similar size; and that phenylalanine, tryptophan and tyrosine all have a generally similar shape. Therefore, based upon these considerations, arginine, lysine and histidine; alanine, glycine and serine; and phenylalanine, tryptophan and tyrosine; are defined herein as conservative amino acid changes or substitutions. In general, conservatively modified variants of a sequence may include one or more conservative amino acid change or substitution.

In making such changes, the hydropathic index of amino acids may be considered. Each amino acid has been assigned a hydropathic index on the basis of their hydrophobicity and charge characteristics, these are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

The importance of the hydropathic amino acid index in conferring interactive biological function on a protein is generally understood in the art (Kyte *et al.*, J. Mol. Biol. (1982) 157(1):105-32, incorporated herein by reference). It is known that certain amino acids may be substituted for other amino acids having a similar hydropathic index or score and still retain a similar biological activity. In making changes based upon the hydropathic index, the substitution of amino acids whose hydropathic indices are within ± 2 is preferred, those which are within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred.

It also is understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity. U.S. Patent 4,554,101 to Hopp,

incorporated herein by reference, states that the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with its immunogenicity and antigenicity, *i.e.* with a biological property of the protein. It is understood that an amino acid can be substituted for
5 another having a similar hydrophilicity value and still obtain a biologically equivalent protein.

As detailed in U.S. Patent 4,554,101, *supra*, the following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0 \pm 1); glutamate (+3.0 \pm 1); serine (+0.3); asparagine (+0.2);
10 glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5 \pm 1); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); tryptophan (-3.4).

In making changes based upon similar hydrophilicity values, the substitution of amino acids whose hydrophilicity values are within \pm 2 is preferred, those which
15 are within \pm 1 are particularly preferred, and those within \pm 0.5 are even more particularly preferred.

While discussion has focused on conservatively modified variant polypeptides and functionally equivalent polypeptides arising from amino acid changes, it will be appreciated that these changes may be effected by alteration of the encoding
20 polynucleotide; taking into consideration also that the genetic code is degenerate and that two or more codons may code for the same amino acid.

28.0 Sequence Modification Techniques

Modifications to sequences, such as MSP sequences, may be made during chemical synthesis of the polymers (either nucleotide or peptide synthesis). It is
25 believed, however, that site-directed mutagenesis of an encoding nucleic acid,

creating a suitably altered polynucleotide sequence is the most cost effective method of generating an altered polynucleotide sequence. Where the MSP protein is desired, then the mutated sequence may be expressed including in culture (*in vitro* or *ex vivo*) or *in vivo*.

5 Site-directed mutagenesis is a technique useful in the preparation of individual peptides, or biologically functional equivalent proteins or peptides, through specific mutagenesis of the underlying DNA. Several methods for site directed mutagenesis are described in U.S. Patent 4,873,192 to Kunkel, incorporated herein by reference and in U.S. Patent 4,351,901 to Ball, incorporated
10 herein by reference. The technique further provides a ready ability to prepare and test sequence variants, for example, incorporating one or more of the foregoing considerations, by introducing one or more nucleotide sequence changes into the DNA. Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences which encode the DNA sequence of the desired
15 mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Typically, a primer of about 14 to about 25 nucleotides in length is preferred, with about 5 to 10 residues on both sides of the junction of the sequence being altered. The primers can be selected by one
20 with ordinary skill in the art based upon information provided herein, including the Sequence Listings and Figures.

 The technique of site-specific mutagenesis is well known in the art as exemplified by publications (Adelman *et al.*, (1983) DNA 2(3)183-193, incorporated
 herein by reference). As will be appreciated, the technique typically employs a
25 phage vector which exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13

phage. These phage are readily commercially available and their use is well known to those skilled in the art. Double stranded plasmids are also routinely employed in site directed mutagenesis which eliminates the step of transferring the gene of interest from a plasmid to a phage. Kits for phage based site directed mutagenesis
5 are commercially available. In addition PCR based methods which may, or may not, involve phage are known in the art and kits for such purposes are commercially available.

10 In certain known techniques, site-directed mutagenesis is performed by first obtaining a single-stranded vector or melting apart the two strands of a double stranded vector which includes within its sequence a DNA sequence which encodes the desired nucleotide, such as MSP. An oligonucleotide primer bearing the desired mutated sequence is prepared, generally synthetically as is known to one of ordinary skill in the art. This primer is then annealed with the single-stranded
15 vector, and subjected to DNA polymerizing enzymes such as Escherichia coli (E. coli) polymerase I Klenow fragment, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform appropriate cells, such as E. coli cells, and clones are selected which include recombinant vectors bearing
20 the mutated sequence arrangement. Various selection methods that increase the percentage of specifically modified clones over wild-type are known and available commercially.

Kalderon *et al.* (1984) report several mutagenic methods which have proved useful in mutating the native LT gene. Specifically, Kalderon *et al.* teach deletion
25 mutations by displacement-loop mutagenesis and by the random insertion of Eco RI linkers into the LT gene. Further, point mutation by deletion-loop mutagenesis is

taught. The reference also teaches screening procedures for determining the success of such mutations. The teachings of Kalderon *et al.* (1984) Virology 139(1)109-137 are incorporated herein by reference.

5 The preparation of sequence variants of the selected gene using site-directed mutagenesis is provided as a method of producing potentially useful nucleic acids and peptides (such as MSP and MSP variants) and is not meant to be limiting as there are other ways in which sequence variants of these nucleotide and peptides may be obtained. For example, recombinant vectors encoding the desired genes may be treated with mutagenic agents to obtain sequence variants for the mutagenesis of plasmid DNA using hydroxylamine or random mutagenesis may be performed using the PCR technique.

10 Sequence analysis of a potentially mutant nucleic acid sequence is carried out by methods known in the art, typically by either Sanger dideoxy sequencing (Sanger *et al.*, PNAS (1977) 74:5363-5467, incorporated herein by reference; U.S. Patent 4,871,929 to Barnes; and U.S. Patent 4,962,020 to Tabor *et al.*, each patent incorporated herein by reference) or automated sequencing (U.S. Patent 5,365,455 to Tibbetts *et al.*, incorporated herein by reference).

15 In addition to the MSP peptidyl compounds described herein, the inventors also contemplate that other sterically similar compounds may be formulated to mimic the key portions of the peptide structure. Such compounds may be used in the same manner as the peptides of the invention and hence are also functional equivalents. The generation of a structural functional equivalent may be achieved by the techniques of modeling and chemical design known to those of skill in the art. It will be understood that all such sterically similar constructs fall within the scope of the present invention.

Livestock include, but are not limited to: horses, work horses, show horses, cattle, sheep, goats, and the like. Pets include, but are not limited to: dogs, cats, horses, birds, and the like.

EXAMPLES

Example 1

Large quantities of sperm (generally >10⁸) are purified using a modification of methods developed by Klass and Hirsh (1981). Adult males are identified using synchronized cultures (Lewis and Flemming, 1995) of *fog-2(q71)*, which are 50% male and 50% female. Mutations in the *fog-2* gene block spermatogenesis in XX animals, transforming them into females, but have no effect on XO animals, which are fertile males (Schedl and Kimble, 1988). Males are separated from females, larvae, and embryos based on size by sieving through NITEX screens of various pore sizes. The populations of males isolated in this way are generally >99% pure. To isolate sperm, males are placed between two PLEXIGLASS plates and smashed in a vice grip (The Home Depot, Inc.). Intact sperm are then purified from the carcasses by filtration through NITEX filters (20 micron pore size) and washed in M9 phosphate buffer (Sulston and Hodgkin, 1988) using several rounds of low speed centrifugation (e.g., 10,000xg) and resuspension. Sperm-conditioned medium (SCM) is prepared by incubating purified sperm in M9 for different periods of time (1-12 h) and subsequently removing the sperm by centrifugation and filtration through a 0.2 micron filter. Microscopic analysis suggests that the sperm are not lysing during the incubation. Polyacrylamide gel electrophoresis (PAGE) further indicates that the sperm are not lysing during the incubation, but that a 14 kDa protein is enriched in SCM (FIG. 2).

REFERENCES

All references, articles, U.S. Patents, Non-U.S. Patents, exhibits and the like referred to herein, including those listed below or attached, are hereby incorporated herein by reference in their entirety.

- 5 Achanzar, W. E., and Ward, S. (1997). A nematode gene required for sperm vesicle fusion. *Journal of Cell Science* 110: 1073-1081.

Albertson, D. G. (1984). Formation of the first cleavage spindle in nematode embryos. *Developmental Biology* 101: 61-72.

- 10 Albertson, D. G., and Thomson, J. N. (1993). Segregation of holocentric chromosomes at meiosis in the nematode, *Caenorhabditis elegans*. *Chromosome Research* 1: 15-26.

Anderson, E., and Albertini, D. F. (1976). Gap junctions between the oocyte and companion follicle cells in the mammalian ovary. *Journal of Cell Biology* 71: 680-686.

- 15 Argon, Y., and Ward, S. (1980). *C. elegans* fertilization-defective mutants with abnormal sperm. *Genetics* 96: 413-433.

Aroian, R. V., Field, C., Pruliere, G., Kenyon, C., and Alberts, B. M. (1997). Isolation of actin-associated proteins from *Caenorhabditis elegans* oocytes and their localization in the early embryo. *EMBO Journal* 16: 1541-1549.

- 20 Aruffo, A., and Seed, B. (1987). Molecular cloning of a CD28 cDNA by a high-efficiency COS cell expression system. *Proceedings of the National Academy of Sciences U S A* 84: 8573-8577.

- 25 Austin, J., and Kimble, J. (1987). *glp-1* is required in the germ line for regulation of the decision between mitosis and meiosis in *C. elegans*. *Cell* 51: 589-599.

Barton, M. K., and Kimble, J. (1990). *fog-1*, a regulatory gene required for specification of spermatogenesis in the germ line of *Caenorhabditis elegans*. Genetics 125: 29-39.

5 Bashir, R., Britton, S., Strachan, T., Keers, S., Vafiadaki, E., Lako, M.,
Richard, I., Marchand, S., Bourg, N., Argov, Z., Sadeh, M., Mahjneh, I., Marconi, G.,
Passos-Bueno, M. R., Moreira, E. de S., Zatz, M., Beckmann, J. S., and Bushby, K.
(1998). A gene related to *Caenorhabditis elegans* spermatogenesis factor *fer-1* is
mutated in limb-girdle muscular dystrophy type 2B. Nature Genetics 20: 37-42.

10 Blaxter, M. (1998). *Caenorhabditis elegans* is a nematode. Science 282: 2041-
2046.

Boxem, M., Srinivasan, D. G., and van den Heuvel, S. (1999). The
Caenorhabditis elegans gene *ncc-1* encodes a cdc2-related kinase required for M
phase in meiotic and mitotic cell divisions, but not for S phase. Development 126:
2227-2239.

15 Brenner, S. (1974). The genetics of *Caenorhabditis elegans*. Genetics 77: 71-
94.

Browning, H., and Strome, S. (1996). A sperm-supplied factor required for
embryogenesis in *C. elegans*. Development 122: 391-404.

20 Bullock, T. L., Roberts, T. M., and Stewart, M. (1996). 2.5 A resolution crystal
structure of the motile major sperm protein (MSP) of *Ascaris suum*. Journal of
Molecular Biology 263: 284-296.

C. elegans Sequencing Consortium. (1998). Genome sequence of the nematode
Caenorhabditis elegans: A platform for investigating biology. Science 282: 2012-
2018.

25 Cadigan, K. M., and Nusse, R. (1997). Wnt signaling: a common theme in
animal development. Genes and Development 11: 3286-3305.

Chase, D., Serafinas, C., Ashcroft, N., Kosinski, M., Longo, D., Ferris, D. K., and Golden, A. (2000). The polo-like kinase PLK-1 is required for nuclear envelope breakdown and the completion of meiosis in *Caenorhabditis elegans*. *Genesis: The Journal of Genetics and Development* 26: 26-41.

- 5 Chen, P. J., Singal, A., Kimble, J., and Ellis, R. E. (2000). A novel member of the Tob family of proteins controls sexual fate in *Caenorhabditis elegans* germ cells. *Developmental Biology* 217: 77-90.

10 Church, D. L., Guan, K.-L., and Lambie, E.J. (1995). Three genes of the MAP kinase cascade, *mek-2*, *mpk-1/sur-1* and *let-60 ras*, are required for meiotic cell cycle progression in *Caenorhabditis elegans*. *Development* 121: 2525-2535.

Clandinin, T. R., DeModena, J. A., and Sternberg, P. W. (1998). Inositol triphosphate mediates a RAS-independent response to LET-23 receptor tyrosine kinase activation in *C. elegans*. *Cell* 92: 523-533.

- 15 Colledge, W. H., Carlton, M. B. L., Udy, G. B., and Evans, M. J. (1994). Disruption of *c-mos* causes parthenogenetic development of unfertilized mouse eggs. *Nature* 370: 65-68.

20 Cordero, M. M., Cornish, T. J., Cotter, R. J., and Lys, I. A. (1995). Sequencing peptides without scanning the reflectron: post-source decay with a curved-field reflectron time-of-flight mass spectrometer. *Rapid Communications in Mass Spectrometry* 9: 1356-1361.

Coulson, A., Huynh, C., Kozono, Y., and Shownkeen, R. (1995). The physical map of the *Caenorhabditis elegans* genome. In: Epstein, H. F., and Shakes, D. C., editors. *Methods in Cell Biology. Caenorhabditis elegans: Modern Biological Analysis of an Organism*. San Diego: Academic Press. p. 533-550.

Cross, D. A., and Smythe, C. (1998). PD98059 prevents establishment of the spindle assembly checkpoint and inhibits the G2-M transition in meiotic but not mitotic cell cycles in *Xenopus*. *Experimental Cell Research* 241: 12-22.

5 Davis, S., Aldrich, T. H., Valenzuela, D. M., Wong, V. V., Furth, M. E., Squinto, S. P., and Yancopoulos, G. D. (1991). The receptor for ciliary neurotrophic factor. *Science* 253: 59-63.

10 Dernburg, A. F., McDonald, K., Moulder, G., Barstead, R., Dresser, M., and Villeneuve, A. M. (1998). Meiotic recombination in *C. elegans* initiates by a conserved mechanism and is dispensable for homologous chromosome synapsis. *Cell* 94: 387-398.

Ellis, R. E., and Kimble, J. (1995). The *fog-3* gene and regulation of cell fate in the germ line of *Caenorhabditis elegans*. *Genetics* 139: 561-577.

15 Ferby, I., Blazquez, M., Palmer, A., Eritja, R., and Nebreda, A. R. (1999). A novel p34(cdc2)-binding and activating protein that is necessary and sufficient to trigger G(2)/M progression in *Xenopus* oocytes. *Genes and Development* 13: 2177-2189.

Ferrell, J. E., Jr. (1999). *Xenopus* oocyte maturation: new lessons from a good egg. *BioEssays* 21: 833-842.

20 Ferrell, J. E., Jr., Wu, M., Gerhart, J. C., and Martin, G. S. (1991). Cell cycle tyrosine phosphorylation of p34cdc2 and a microtubule-associated protein kinase homolog in *Xenopus* oocytes and eggs. *Molecular and Cellular Biology* 11: 1965-1971.

25 Fire, A., Xu, S., Montgomery, M. K., Kostas, S. A., Driver, S. E, and Mello, C. C. (1998). Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 391: 806-811.

Fisher, D. L., Brassac, T., Galas, S., and Doree, M. (1999). Dissociation of MAP kinase activation and MPF activation in hormone-stimulated maturation of *Xenopus* oocytes. *Development* 126: 4537-4546.

Francis, R., Barton, M. K., Kimble, J., and Schedl, T. (1995). *gld-1*, a tumor
5 suppressor gene required for oocyte development in *Caenorhabditis elegans*.
Genetics 139: 579-606.

Gavrilets, G., *Nature* 403, 886 (2000).

Godeau, J. F., Schorderet-Slatkine, S., Hubert, P., and Baulieu, E. E. (1978).
Induction of maturation in *Xenopus laevis* oocytes by a steroid linked to a polymer.
10 *Proceedings of the National Academy of Sciences USA* 75: 2353-2357.

Gotoh, Y., Masuyama, N., Dell, K., Shirakabe, K., and Nishida, E. (1995).
Initiation of *Xenopus* oocyte maturation by activation of the mitogen-activated
protein kinase cascade. *Journal of Biological Chemistry* 270: 25898-25904.

Gotoh, Y., Moriyama, K., Matsuda, S., Okumura, E., Kishimoto, T.,
15 Kawasaki, H., Suzuki, K., Yahara, I., Sakai, H., and Nishida, E. (1991). *Xenopus* M
phase MAP kinase: isolation of its cDNA and activation by MPF. *EMBO Journal* 10
:2661-2668.

Grant, B., and Hirsh, D. (1999). Receptor-mediated endocytosis in the
Caenorhabditis elegans oocyte. *Molecular Biology of the Cell* 10: 4311-4326.

20 Greenstein, D., Hird, S., Plasterk, R. H. A., Andachi, Y., Kohara, Y., Wang,
B., Finney, M., and Ruvkun, G. (1994). Targeted mutations in the *Caenorhabditis*
elegans POU homeo box gene *ceh-18* cause defects in oocyte cell cycle arrest, gonad
migration, and epidermal differentiation. *Genes and Development* 8: 1935-1948.

Greenstein, D., unpublished results.

Gross, S. D., Schwab, M. S., Taieb, F. E., Lewellyn, A. L., Qian, Y. W., and Maller, J. L. (2000). The critical role of the MAP kinase pathway in meiosis II in *Xenopus oocytes* is mediated by p90(Rsk). *Current Biology* 10: 430-438.

5 Haaf, A., Butler, P. J., Kent, H. M., Fearnley, I. M., Roberts, T. M., Neuhaus, D., and Stewart, M. (1996). The motile major sperm protein (MSP) from *Ascaris suum* is a symmetric dimer in solution. *Journal of Molecular Biology* 260: 251-260.

Haccard, O., Lewellyn, A., Hartley, R. S., Erikson, E., and Maller, J. L. (1995). Induction of *Xenopus oocyte* meiotic maturation by MAP kinase. *Developmental Biology* 168: 677-682.

10 Hall, D. H., Winfrey, V. P., Blaeuer, G., Hoffman, L. H., Furuta, T., Rose, K. L., Hobert, O., and Greenstein, D. (1999). Ultrastructural features of the adult hermaphrodite gonad of *Caenorhabditis elegans*: relations between the germ line and soma. *Developmental Biology* 212: 101-123.

15 Hashimoto, N., Watanabe, N., Furuta, Y., Tamemoto, H., Sagata, N., Yokoyama, M., Okazaki, K., Nagayoshi, M., Takeda, N., Ikawa, Y., and Aizawa, S. (1994). Parthenogenetic activation of oocytes in *c-mos* deficient mice. *Nature* 370: 68-71.

20 Hill, D. P., Shakes, D. C., Ward, S., and Strome, S. (1989). A sperm-supplied product essential for initiation of normal embryogenesis in *Caenorhabditis elegans* is encoded by the paternal-effect embryonic-lethal gene, *spe-11*. *Developmental Biology* 136: 154-166.

Hill, K. L. and L'Hernault, S. W., submitted.

Hirsh, D., Oppenheim, D., and Klass, M. (1976). Development of the reproductive system of *Caenorhabditis elegans*. *Developmental Biology* 49: 200-219.

25 Holst, P. A., and Phemister, R. D. (1971). The prenatal development of the dog: Preimplantation events. *Biology of Reproduction* 5: 194-206.

Hooper, N. M., and Bashir, A. (1991). Glycosyl-phosphatidylinositol-anchored membrane proteins can be distinguished from transmembrane polypeptide-anchored proteins by differential solubilization and temperature-induced phase separation in Triton X-114. *Biochemical Journal* 280: 745-751.

5 Huang, W., Kessler, D. S., and Erikson, R. L. (1995). Biochemical and biological analysis of Mek1 phosphorylation site mutants. *Molecular Biology of the Cell* 6: 237-245.

Hubbard, E. J. A., and Greenstein, D. (2000). The *C. elegans* gonad: a test tube for cell and developmental biology. *Developmental Dynamics* 218: 2-22.

10 Hyttel, P., Farstad, W., Mondain-Monval, M., Bakke Lajord, K., and Smith, A. J. (1990). Structural aspects of oocyte maturation in the blue fox (*Alopex lagopus*). *Anatomy and Embryology* 181, 325-331.

15 Ishikawa, K., Hanaoka, Y., Kondo, Y., and Imai, K. (1977). Primary action of steroid hormone at the surface of amphibian oocyte in the induction of germinal vesicle breakdown. *Molecular and Cellular Endocrinology* 9: 91-100.

Italiano, J. E., Jr., Roberts, T. M., Stewart, M., and Fontana, C. A. (1982). Reconstitution *in vitro* of the motile apparatus from the amoeboid sperm of *Ascaris* shows that filament assembly and bundling move membranes. *Cell* 84: 105-114.

20 Jansen, G., Hazendonk, E., Thijssen, K. L., and Plasterk, R. H. (1997). Reverse genetics by chemical mutagenesis in *Caenorhabditis elegans*. *Nature Genetics* 17: 119-121.

25 Kagiwada, S., Hosaka, K., Murata, M., Nikawa, J., and Takatsuki, A. (1998). The *Saccharomyces cerevisiae* SCS2 gene product, a homolog of a synaptobrevin-associated protein, is an integral membrane protein of the endoplasmic reticulum and is required for inositol metabolism. *Journal of Bacteriology* 180: 1700-1708.

Kanatani, H., Shirai, H., Nakanishi, K., and Kurokawa, T. (1969). Isolation and identification on meiosis inducing substance in starfish *Asterias amurensis*. Nature 221: 273-274.

5 Kaufmann, R., Spengler, B., and Lutzenkirchen, F. (1993). Mass spectrometric sequencing of linear peptides by product-ion analysis in a reflectron time-of-flight mass spectrometer using matrix-assisted laser desorption ionization. Rapid Communications in Mass Spectrometry 7: 902-910.

10 Kimble, J., and Hirsh, D. (1979). The postembryonic cell lineages of the hermaphrodite and male gonads in *Caenorhabditis elegans*. Developmental Biology 70: 396-417.

Kirby, C., Kusch, M., and Kempfues, K. (1990). Mutations in the *par* genes of *Caenorhabditis elegans* affect cytoplasmic reorganization during the first cell cycle. Developmental Biology 142: 203-215.

15 Klass, M. R., and Hirsh, D. (1981). Sperm isolation and biochemical analysis of the major sperm protein from *C. elegans*. Developmental Biology 84: 299-312.

Kosako, H., Gotoh, Y., and Nishida, E. (1994). Requirement for the MAP kinase kinase/MAP kinase cascade in *Xenopus* oocyte maturation. EMBO Journal 3: 2131-2138.

20 LaMunyon, C. W., and Ward, S. (1994). Assessing the viability of mutant and manipulated sperm by artificial insemination of *Caenorhabditis elegans*. Genetics 138: 689-692.

Laurent, F., Labesse, G., and de Wit, P. (2000). Molecular cloning and partial characterization of a plant VAP33 homologue with a major sperm protein domain. Biochemical and Biophysical Research Communications 270: 286-292.

25 Lee, M.-H. and Schedl, T., unpublished results.

Lewis, J. A., and Flemming, J. T. (1995). Basic culture methods. In: Epstein, H. F., and Shakes, D. C., editors. *Methods in Cell Biology. Caenorhabditis elegans: Modern Biological Analysis of an Organism*. San Diego: Academic Press. p. 3-29.

L'Hernault, S. W. (1997). Spermatogenesis. In: Riddle, D. L., Blumenthal, T., Meyer, B. J., and Priess, J. R., editors. *C. elegans II*. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press. p. 271-294.

L'Hernault, S. W., and Roberts, T. M. (1995). Cell biology of nematode sperm. In: Epstein, H. F., and Shakes, D. C., editors. *Methods in Cell Biology. Caenorhabditis elegans: Modern Biological Analysis of an Organism*. San Diego: Academic Press. p. 273-301.

L'Hernault, S. W., Arduengo, P. M., *J. Cell Biol.* 119, 55 (1992).

Lin, R.J., Kao, H. Y., Ordentlich, P., and Evans, R. M. (1998). The transcriptional basis of steroid physiology. *Cold Spring Harbor Symposium on Quantitative Biology* 63: 577-585.

Liu, X., Kim, C. N., Yang, J., Ronald Jemmerson, R., and Wang, X. (1996). Induction of apoptotic program in cell-free extracts: requirement for dATP and cytochrome c. *Cell* 86: 147-157.

MacMorris, M., Spieth, J., Madej, C., Lea, K., and Blumenthal, T. (1994). Analysis of the VPE sequences in the *Caenorhabditis elegans vit-2* promoter with extrachromosomal tandem array-containing transgenic strains. *Molecular and Cellular Biology* 14: 484-491.

Masui, Y., and Clarke, H. J. (1979). Oocyte maturation. *International Review of Cytology* 57: 185-282.

Masui, Y., and Markert, C. L. (1971). Cytoplasmic control of nuclear behavior during meiotic maturation of frog oocytes. *Journal of Experimental Zoology* 177: 129-145.

McCarter, J., Bartlett, B., Dang, T., and Schedl, T. (1997). Soma-germ cell interactions in *Caenorhabditis elegans*: Multiple events in germline development require the somatic sheath and spermathecal lineages. *Developmental Biology* 181: 121-143.

- 5 McCarter, J., Bartlett, B., Dang, T., and Schedl, T. (1999). On the control of oocyte meiotic maturation and ovulation in *C. elegans*. *Developmental Biology* 205: 111-128.

10 Mendez, R., Hake, L. E., Andresson, T., Littlepage, L. E., Ruderman, J. V., and Richter, J. D. (2000). Phosphorylation of CPE binding factor by Eg2 regulates translation of *c-mos* mRNA. *Nature* 404: 302-307.

15 Miller, M. A., Nguyen V. Q., Lee, M., Kosinski, M., Schedl, M., Caprioli, R. M., and Greenstein D. (2001) A Sperm Cytoskeletal Protein That Signals Oocyte Meiotic Maturation and Ovulation. *Science* 2001 291: 2144-2147.

20 Morgan, D. O. (1997). Cyclin-dependent kinases: engines, clocks, and microprocessors. *Annual Review of Cell and Developmental Biology* 13: 261-291.

25 Muhlrad, D., Hunter, R., and Parker, R. (1992). A rapid method for localized mutagenesis of yeast genes. *Yeast* 8: 79-82.

 Mukherjee, S., Ghosh, R. N., and Maxfield, F. R. (1997). Endocytosis. *Physiology Reviews* 77: 759-803.

 Munson, P. J., and Rodbard, D. (1980). Ligand: a versatile computerized approach for characterization of ligand-binding systems. *Analytical Biochemistry* 107: 220-239.

 Myers, C. D., Goh, P.-Y., Allen, T. St. C., Bucher, E. A., and Bogaert, T. (1996). Developmental genetic analysis of Troponin T mutations in striated and nonstriated muscle cells of *Caenorhabditis elegans*. *Journal of Cell Biology* 132: 1061-1077.

Nance, J., Minniti, A. N., Sadler, C., and Ward, S. (1999). *spe-12* encodes a sperm cell surface protein that promotes spermiogenesis in *Caenorhabditis elegans*. *Genetics* 152: 209-220.

5 Nelson, G. A., Roberts, T. M., and Ward, S. (1982). *C. elegans* spermatozoan locomotion: Amoeboid movement with almost no actin. *Journal of Cell Biology* 92: 121-131.

Nelson, G. A., and Ward, S. (1980). Vesicle fusion, pseudopod extension and amoeboid motility are induced in nematode spermatids by the ionophore monensin. *Cell* 19: 457-464.

10 Palmer, A. and Nebreda, A. R., *Prog. Cell Cycle Res.* 4, 131 (2000).

Pandey, A., and Mann, M. (2000). Proteomics to study genes and genomes. *Nature* 405: 837-846.

15 Pavalko, F. M., and Roberts, T. M. (1989). Posttranslational insertion of a membrane protein on *Caenorhabditis elegans* sperm occurs without de novo protein synthesis. *Journal of Cellular Biochemistry* 41: 57-70.

Podbilewicz, B. (1996). ADM-1, a protein with metalloprotease- and disintegrin-like domains, is expressed in syncytial organs, sperm, and sheath cells of sensory organs in *Caenorhabditis elegans*. *Molecular Biology of the Cell* 7: 1877-1893.

20 Posada, J., and Cooper, J. A. (1992). Requirements for phosphorylation of MAP kinase during meiosis in *Xenopus* oocytes. *Science* 255: 212-215.

Qian, Y. W., Erikson, E., and Maller, J. L. (1999). Mitotic effects of a constitutively active mutant of the *Xenopus* polo-like kinase Plx1. *Molecular and Cellular Biology* 19: 8625-8632.

25 Ramalho-Santos, J., *et al.*, *Dev. Biol.* 223, 54 (2000).

Resing, K. A., Mansour, S. J., Hermann, A. S., Johnson, R. S., Candia, J. M., Fukasawa, K., Vande Woude, G. F., and Ahn, N. G. (1995). Determination of v-Mos-catalyzed phosphorylation sites and autophosphorylation sites on MAP kinase by ESI/MS. *Biochemistry* 34: 2610-2620.

5 Roberts, T. M. (1983). Crawling *C. elegans* spermatozoa contact the substrate only by their pseudopods and contain 2-nm filaments. *Cell Motility* 3: 333-347.

Roberts, T. M., Pavalko, F. M., and Ward, S. (1986). Membrane and cytoplasmic proteins are transported in the same organelle complex during nematode spermatogenesis. *Journal of Cell Biology* 102: 1787-1796.

10 Roberts, T. M., and Stewart, M. (1995). Nematode sperm locomotion. *Current Opinion in Cell Biology* 7: 13-17.

Roberts, T. M., and Stewart, M. (2000). Acting like actin. The dynamics of the nematode major sperm protein (msp) cytoskeleton indicate a push-pull mechanism for amoeboid cell motility. *Journal of Cell Biology* 149: 7-12.

15 Roberts, T. M., and Ward, S. (1982a). Membrane flow during nematode spermiogenesis. *Journal of Cell Biology* 92: 113-120.

Roberts, T. M., and Ward, S. (1982b). Centripetal flow of pseudopodial surface components could propel the amoeboid movement of *C. elegans* spermatozoa. *Journal of Cell Biology* 92: 132-138.

20 Rose, K. L., Winfrey, V. P., Hoffman, L. H., Hall, D. H., Furuta, T., and Greenstein D. (1997). The POU gene *ceh-18* promotes gonadal sheath cell differentiation and function required for meiotic maturation and ovulation in *Caenorhabditis elegans*. *Developmental Biology* 192: 59-77.

25 Rutledge, E., Bianchi, L., Christensen, M., Morrison, R., Broslat, A., Beld, A., George Jr., A. L., Greenstein, D., and Strange, K. (2000). CLH-3: a *C. elegans* ClC-2

chloride channel orthologue regulated by cell cycle events and involved in soma-germline intercellular signaling. Submitted.

Sagata, N. (1997). What does Mos do in oocytes and somatic cells? *BioEssays* 19: 13-21.

5 Sagata, N., Oskarsson, M., Copeland, T., Brumbaugh, J., and Vande Woude, G. F. (1988). Function of *c-mos* proto-oncogene product in meiotic maturation in *Xenopus* oocytes. *Nature* 335:519-525.

10 Sagata, N., Watanabe, N., Vande Woude, G. F., and Ikawa, Y. (1989). The *c-mos* proto-oncogene product is a cytostatic factor responsible for meiotic arrest in vertebrate eggs. *Nature* 342: 512-518.

Schagger, H., and von Jagow, G. (1987). Tricine-sodium dodecyl sulfate-polyacrylamide gel electrophoresis for the separation of proteins in the range from 1 to 100 kDa. *Analytical Biochemistry* 166: 368-379.

15 Schedl, T., and Kimble, J. (1988). *fog-2*, a germ-line-specific sex determination gene required for hermaphrodite spermatogenesis in *C. elegans*. *Genetics* 119: 43-61.

Schulz, J. R., *et al.*, *J. Biol. Chem.* 273, 24355 (1998).

20 Seed, B., and Aruffo, A. (1987). Molecular cloning of the CD2 antigen, the T-cell erythrocyte receptor, by a rapid immunoselection procedure. *Proceedings of the National Academy of Sciences U S A* 84: 3365-3369.

Shakes, D. C., and Ward, S. (1989). Initiation of spermiogenesis in *C. elegans*: A pharmacological and genetic analysis. *Developmental Biology* 134: 189-200.

Sheets, M. D., Wu, M., and Wickens, M. (1995). Polyadenylation of *c-mos* mRNA as a control point in *Xenopus* meiotic maturation. *Nature* 374: 511-516.

25 Skehel, P. A., Armitage, B. A., Bartsch, D., Hu, Y., Kaang, B. K., Siegelbaum, S. A., Kandel, E. R., and Martin, K. C. (1995). Proteins functioning in synaptic

transmission at the sensory to motor synapse of *Aplysia*. *Neuropharmacology* 34: 1379-1385.

Skehel, P. A. *et al.*, *Science* **269**, 1580 (1995).

Smith, H. E., and Ward, S. (1998). Identification of protein-protein
5 interactions of the major sperm protein (MSP) of *Caenorhabditis elegans*. *Journal of Molecular Biology* 279: 605-619.

Smith, L. D., and Ecker, R. E. (1969). Role of the oocyte nucleus in physiological maturation in *Rana pipiens*. *Developmental Biology* 19: 281-309.

Smith, L. D., and Ecker, R. E. (1971). The interaction of steroids with *Rana*
10 *pipiens* oocytes in the induction of maturation. *Developmental Biology* 25: 232-247.

Soussan, L. *et al.*, *J. Cell Biol.* **146**, 301 (1999).

Spector, D. L., Goldman, R. D., and Leinwand, L. A. (1998). Plasma
membrane isolation using the cationic colloidal silica isolation technique. In *Culture and Biochemical Analysis of Cells*. K. Janssen, editor. p. 35.1-35.14. Cold Spring
15 Harbor Laboratory Press: Plainview, NY.

Starck, J., Gibert, M.-A., Brun, J., and Bosch C. (1983), Ribosomal RNA
synthesis and processing during oogenesis of the free living nematode *C. elegans*.
Comparative Biochemistry and Physiology 75B: 575-580.

Stebbins-Boaz, B., Hake, L. E., and Richter, J. D. (1996). CPEB controls the
20 cytoplasmic polyadenylation of cyclin, Cdk2 and *c-mos* mRNAs and is necessary for
oocyte maturation in *Xenopus*. *EMBO Journal* 5: 2582-2592.

Strome, S. (1986). Fluorescence visualization of the distribution of
microfilaments in gonads and early embryos of the nematode *C. elegans*. *Journal of Cell Biology* 103: 2241-2252.

Sulston, J., and Hodgkin, J. (1988). Methods. In The Nematode *Caenorhabditis elegans*. W. B. Wood, editor, p. 587-606. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

5 Swofford, D., "PAUP: Phylogenetic analysis using parsimony." Illinois National History Survey, Champaign (1993).

Vacquier, V. D., *Science* 281, 1995 (1998).

Veenstra, J. A. (2000). Mono- and dibasic proteolytic cleavage sites in insect neuroendocrine peptide precursors. *Archives of Insect Biochemistry and Physiology* 43: 49-63.

096300E90E950
10
15
20
25

10 Ward, S. (1986). Asymmetric localization of gene products during the development of *C. elegans* spermatozoa. In Gametogenesis and the Early Embryo. 44th Symposium of the Society for Developmental Biology. Gall, J. G., (ed). p.55-75. Alan R. Liss, NY.

15 Ward, S., Argon, Y., and Nelson, G. A. (1981). Sperm morphogenesis in wild-type and fertilization-defective mutants of *C. elegans*. *Journal of Cell Biology* 91: 26-44.

20 Ward, S., Burke, D. J., Sulston, J. E., Coulson, A. R., Albertson, D. G., Ammons, D., Klass, M., and Hogan, E. (1988). Genomic organization of major sperm protein genes and pseudogenes in the nematode *Caenorhabditis elegans*. *Journal of Molecular Biology* 199, 1-13.

Ward, S., and Carrel, J. S. (1979). Fertilization and sperm competition in the nematode *Caenorhabditis elegans*. *Developmental Biology* 73: 304-321.

Ward, S., Hogan, E., and Nelson, G. A. (1983). The initiation of spermiogenesis in the nematode *C. elegans*. *Developmental Biology* 98: 70-79.

25 Ward, S., and Klass, M. (1982). The location of the major protein in *C. elegans* sperm and spermatocytes. *Developmental Biology* 92: 203-208.

Ward, S., and Miwa, J. (1978). Characterization of temperature-sensitive, fertilization-defective mutants of the nematode *C. elegans*. *Genetics* 88: 285-303.

Ward, S., Roberts, T. M., Strome, S., Pavalko, F. M., and Hogan, E. (1986). Monoclonal antibodies that recognize a polypeptide antigenic determinant shared
5 by multiple *C. elegans* sperm-specific proteins. *Journal of Cell Biology* 102: 1778-1786.

Wimalawansa, S. J. (1995). Purification and biochemical characterization of Neuropeptide Y₂ receptor. *The Journal of Biological Chemistry* 270: 18523-18530.

Wolf, N., Hirsh, D., and McIntosh, J. R. (1978). Spermatogenesis in males of
10 the free-living nematode, *C. elegans*. *Journal of Ultrastructure Research* 63: 155-169.

Yasunaga, S., Grati, M., Cohen-Salmon, M., El-Amraoui, A., Mustapha, M., Salem, N., El-Zir, E., Loiselet, J., and Petit, C. (1999). A mutation in OTOF, encoding otoferlin, a FER-1-like protein, causes DFNB9, a nonsyndromic form of
15 deafness. *Nature Genetics* 21:363-369.

Yew, N., Mellini, M. L., and Vande Woude, G. F. (1992). Meiotic initiation by the mos protein in *Xenopus*. *Nature* 355: 649-652.

Yung, Y., *et al.*, *FEBS Lett.* 408, 292 (1997).

4. To prepare SCM, purified sperm (26) were incubated in M9 buffer (~5 x 10⁷
20 sperm/ml) for 1-16 hrs at 20°C. Sperm were removed by centrifugation for 5 min at 14,000 rpm in an Eppendorf microcentrifuge (model 5415C) and filtration through a 0.22 µm cellulose acetate filter (Costar). Samples (~50 pl) were microinjected into the uterus of *fog-2(q71)* (3) adult females (30 hrs post-L4 at 20°C). Following injection, females were anesthetized for 20 minutes with 0.1% tricaine/0.01%
25 tetramisole (32) in M9. Oocyte maturation and sheath cell contraction rates were monitored by time-lapse video microscopy (1) for 70 minutes.

5. SCM or sperm lysates (by vortexing with glass beads) were fractionated on C₄ and C₁₈ columns (Vydac) using an acetonitrile gradient (0-100%) mobile phase. TFA (0.1%) was added to the mobile phase to sharpen peaks by ion pairing. Absorbance peaks (214 nm) were collected manually, dialyzed against M9, and
5 bioassayed. Active fractions were analyzed by MALDI-TOF mass spectrometry using internal molecular weight standards (insulin, cytochrome C and myoglobin).

6. Post source decay mass spectrometry (33) of a 1960 Da peptide, generated by tryptic digestion of the active fraction, yielded the sequence IVFNAPYDDKHTYHIK, which matched MSP.

10 9. His-tagged MSP-77, MSP-38, and MSP(1-106) were expressed in *Escherichia coli* M15[pRep4] (Qiagen) and purified under native conditions by Ni-NTA affinity chromatography (>99% pure by SDS-PAGE and mass spectrometry). The extinction coefficient ϵ (275 nm) of MSP was estimated by amino acid analysis of a purified His-tagged MSP-77 standard. MSP concentrations were determined by
15 amino acid hydrolysis, SDS-PAGE, and spectrophotometrically using ϵ (275 nm) = $3.29 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$.

20 10. Anti-MSP (26) or control EMB-30 antibodies (34) were injected (~40 $\mu\text{g/ml}$) into adult wild-type N2 hermaphrodites (24 hr post-L4 at 20°C). The injected animals were cultured individually with food for a 3 hr time period and total ovulations were determined. The effect of antibody injection on oocyte maturation was determined by time lapse video microscopy. A two-sample t-test was used to compare results of MSP injections with controls.

12. The C-terminal MSP peptide (EWFQGDGMVRRKNLPIEYNP) was prepared by solid-phase synthesis and purified by HPLC (Research Genetics).

15. Diphosphorylated MAP kinase was detected in dissected and fixed (3% paraformaldehyde) gonadal preparations using indirect immunofluorescence with the antibody MAPK-YT (35) (Sigma). In *C. elegans* preparations, MAPK-YT only recognizes *mpk-1* map kinase gene products (36). Gonads were stained 8, 40, or 50 min post MSP injection. Activated MAP kinase was detected 40 and 50 min post-injection but was not detectable 8 min post injection.

22. Phylogenetic analyses were performed using maximum parsimony and neighbor-joining methods. Amino acid sequences from MSP and MSP-like domains of several representative VAPs were used in the analyses. For parsimony, the heuristic search option of PAUP* 3.1 (37) was used for tree construction, with 200 random order taxon addition replicates and tree bisection and reconnection branch swapping. Bacterial PapD, which is structurally related to MSP, was used as the outgroup. The "protpars" matrix of PAUP* 3.1 was used to weigh amino acid substitutions. To obtain bootstrap values, 100 bootstrap replicates were performed using simple taxon addition with tree bisection and reconnection branch swapping.

The present invention is not limited by mechanism or theory. Although there have been described general and specific embodiments of the invention herein, these embodiments do not limit the scope of the invention except as set forth in the Claims below.